

QUANTIZATION BY COCHAIN TWISTS AND NONASSOCIATIVE DIFFERENTIALS

E.J. BEGGS & S. MAJID

ABSTRACT. We show that several standard associative quantizations in mathematical physics can be expressed as cochain module-algebra twists in the spirit of Moyal products at least to $O(\hbar^3)$, but to achieve this we twist not by a 2-cocycle but by a 2-cochain. This implies a hidden nonassociativity not visible in the algebra itself but present in its deeper noncommutative differential geometry, a phenomenon first seen in our previous work on semiclassicalisation of differential structures. The quantisations are induced by a classical group covariance and include: enveloping algebras $U_\hbar(\mathfrak{g})$ as quantisations of $\mathfrak{g}^*$, a Fedosov-type quantisation of the sphere S^2 under a Lorentz group covariance, the Mackey quantisation of homogeneous spaces, and the standard quantum groups $\mathbb{C}_q[G]$. We also consider the differential quantisation of $\mathbb{R}^n$ for a given symplectic connection as part of our semiclassical analysis and we outline a proposal for the Dirac operator.

1. INTRODUCTION

This paper is a sequel to [BM1] in which we studied algebras that were associative to the required order in a deformation parameter $\hbar$ but allowed the possibility that the exterior algebra in noncommutative geometry could be nonassociative to that order. We showed that this was *necessary* for the standard quantum groups $\mathbb{C}_q[G]$, i.e. these associative algebras admit no associative exterior algebra of classical dimensions that is bicovariant. Nonassociative calculi were, however, possible by use of Drinfeld's twisting [D2] applied in the category of (super)coquasiHopf algebras. In the present work we provide many more examples using not the quasi-Hopf algebra theory itself but a 'module algebra' twist theory in which any algebra in the category of modules covariant under the a classical (or quantum) group is also twisted. Such methods have been used to obtain nonassociative algebras [AM1] as well as associative ones [DGM]. That one obtains differential calculi as well on such algebras is explored in general terms in [AM2].

We show now that this setting also allows to obtain associative algebras and induced differential calculi for some very standard and *not*-quantum-group-related quantizations, but with a similar price to pay. Thus, we use Hopf algebra methods but apply them to classical situations, notably to coadjoint spaces $\mathfrak{g}^*$ and their quantisation by the enveloping algebra $U_\hbar(\mathfrak{g})$. Clearly this and other 'noncommutative coordinate algebras' that we consider are perfectly associative so it is some surprise, and the main result of the present paper, that their natural induced noncommutative differential calculus is again nonassociative. In the case of $U_\hbar(\mathfrak{g})$ we show (Theorem 5.1.2) that any calculus which is translation and $\mathfrak{g}$ -covariant and has classical dimensions must be nonassociative. This is analogous to the result in [BM1] for quantum groups but now for classical enveloping algebras (and the proof is similar). In this way we confirm and provide major new examples of the general analysis in [BM1]. We particularly analyse the semiclassical level of these results in terms of Poisson and symplectic geometry followed by the next-to-semiclassical order.

An outline of the paper is as follows. In Section 2 we describe the general algebraic twisting theory that we shall use. Section 3 then describes the special case that will be used for all our examples,

namely a method of quantisation induced by a classical symmetry and a cochain. Thus we begin with a *classical* manifold M with a classical Lie algebra symmetry group $\mathcal{L} \subseteq \text{diff}(M)$. As Hopf algebra we take $H = U(\mathcal{L})$ the enveloping algebra. Then the scheme is that any suitable element $F \in H \otimes H$ (a cochain) induces a quantisation of M . We semiclassicalise this theory and see how Poisson-compatible (pre)connections in the sense of [BM1] arise out of the choice of F and $\mathcal{L}$. The choice of the latter covariance Lie algebra determines what kind of connections or preconnections can arise by the cochain twisting construction and hence what structures the quantisation respects. We also briefly discuss the inverse problem of obtaining a cochain F and hence a quantisation given a symplectic form and symplectic connection on M . Section 3.2 analyses the situation for $M = \mathbb{R}^{2n}$ with its standard symplectic structure and general symplectic connection.

Sections 4,5,6,7 then turn to the main examples of the paper. These examples are all constructed by a second order or in some case third order analysis, i.e. we obtain the required cochain at least up to and including $\hbar^2$ terms. This will already be a substantial amount of work and is enough to expose the main phenonema. Moreover, the existence of a cochain to all orders is not really in doubt in view of the Kontsevitch universal quantisation theorem (our cochain amounts to choosing a natural ‘lifting’ of that); our results constitute a natural choice at low order and suggest that a natural choice should be possible to all further orders.

We start these examples with Section 4 in which the sphere S^2 has a natural cochain F for covariance Lie algebra $\mathcal{L} = \mathfrak{so}(1,3)$. The action of the Lorentz group that we use is the one on the ‘sphere at infinity’ in 4-dimensional Minkowski space. We show that one obtains an associative quantisation of the sphere at least to $O(\hbar^3)$ and that this coincides with the Fedosov quantisation to this order for the standard Levi-Civita connection on the sphere (which is symplectic).

Section 5 is the main example of interest in the paper. We show that the classical enveloping algebra $U_\hbar(\mathfrak{g})$ viewed as a quantisation of $S(\mathfrak{g}) = \mathbb{C}[\mathfrak{g}^*]$ (functions on $\mathfrak{g}^*$ with its Kirillov-Kostant bracket) can be viewed as a module-algebra cochain twist and that this quantizes a canonical covariant preconnection in the Poisson geometry of $\mathfrak{g}^*$ (we show that this is in fact the only such preconnection for all simple $\mathfrak{g}$ other than $\mathfrak{sl}_n$, $n > 2$ and even there it is the natural choice). The background covariance we use is $\mathcal{L} = \mathfrak{g} \ltimes \mathfrak{g}^*$ and we find a suitable F as a powerseries to $O(\hbar^3)$ and find that it is essentially unique to this level when we demand a further condition (Section 5.4) whereby $S(\mathfrak{g}^*) = \mathbb{C}[\mathfrak{g}] \subset U(\mathcal{L})$ twists into a local version of the *group* coordinate algebra $\mathbb{C}[G]$ (see below). In effect, we require that F implements the Campbell-Baker-Hausdorff formula by conjugation in addition to its other properties. In Section 5.5 we discuss the Duflo map in this context and argue that the reduced form of F should be the coboundary of the Duflo operator (and hence known to all orders). Although our specific universal F is only found to $O(\hbar^3)$ it seems likely that these various features should extend and characterise it completely. This would be a topic for further work beyond our methods here. In Section 5.6 we demonstrate the theory on $\mathfrak{g} = \mathbb{R} \rtimes \mathbb{R} = \mathfrak{b}_+$ the solvable Lie algebra in 2-dimensions. A version of its enveloping algebra has been proposed as ‘noncommutative spacetime’[MR] and we exhibit a (non-unique) F explicitly to $O(\hbar^4)$ in this case.

Section 6 completes our trio of conventional examples with the Mackey quantisation $C^\infty(N) \rtimes U_\hbar(g)$ as a cochain quantisation of $C^\infty(N) \otimes S(g) \subseteq C^\infty(N \times g^*)$. This extends the model in Section 5 but we need an extended cocycle and covariance Lie algebra $\mathcal{L} = \mathfrak{g} \ltimes \mathfrak{g}^* \oplus \mathfrak{g}$ in order to achieve this. Section 6.3 includes the case $C^\infty(G) \rtimes U_\hbar(g)$ as a quantisation of $T^*G = G \times \mathfrak{g}^*$, where g is the Lie algebra of a Lie group G .

We follow these with the more technical example Section 7 from quantum group theory, which is simply Drinfeld’s theory for quantum groups $\mathcal{C}_q[G]$ reworked as a cochain twist. Here $\mathcal{L} = \mathfrak{g} \oplus \mathfrak{g}^{\text{op}}$ acting from the left and right and F, Φ are built from Drinfeld’s ones relating to the KZ-equations.

This example is not fundamentally new but provides the role model for our view of the more conventional quantisations in the paper, so is included for completeness.

Section 8 turns to the hidden nonassociativity that we have identified in the associative quantum algebras above. The most important of the many implications resulting from the cochain twist is in Section 8.2, namely the corresponding differential calculi. Because in our examples F is not a cocycle, the exterior algebra obtained likewise by twisting is not necessarily associative, and we show that indeed it is not for our various examples. We describe the nonassociative differentials for each of our examples to order $O(\hbar^2)$. One example is a more covariant but nonassociative differential calculus for the non-commutative spacetime in [MR]. Section 8.3 shows how the same philosophy can be used to construct Dirac operators in the sense of generalised ‘spectral triples’. The slight generalisation beyond the axioms in [C] reflects the nonassociativity. We show that such deformations are isospectral, a point of view consistent with other approaches such as [DLSSV].

It is also true that under the cochain quantisation scheme the original covariance becomes a (quasi)quantum group H_F covariance, which we describe to $\hbar^2$ in Section 8.1 for each of our examples. In the case of $U_\hbar(g)$ it appears by accident to be a usual (not quasi) quantum group and to be a local version of the quantum double $D(U(g)) = U(g) \ltimes \mathbb{C}[G]$, which is known to be a covariance quantum group of $U(g)$. When $U_\hbar(su_2)$ is viewed as noncommutative $\mathbb{R}^3$ (the so-called universal fuzzy-sphere), for example, the quantum double plays the role of quantum Euclidean group[BaMa] motivated from 2+1 quantum gravity. In this case the curvature of the canonical preconnection or the fact that F cannot be taken to be a cocycle represents an anomaly in this quantisation of $\mathbb{R}^3$. The associativity obstruction in this case can in fact be resolved by adjoining an extra ‘time’ variable and has been proposed [M4] as an origin of time in noncommutative differential geometry.

Let us say finally that we work in a deformation-theoretic setting with all deformed expressions given by power-series in a parameter $\hbar$ and otherwise over $\mathbb{C}$; all constructions can be formulated more (co)algebraically over any field using comodules which would, however, be less familiar to most readers. In the main ‘examples’ sections we work only to lower degrees in $\hbar$ for which purposes one may regard $\hbar$ as a real parameter with the deformed product of smooth functions assumed to have these first terms in an expansion. The authors would like to thank F.W. Clarke for his assistance with some of the MATHEMATICA calculations underlying the paper and Y. Bazlov for drawing our attention to the Duflo map.

2. PRELIMINARIES: MODULE-ALGEBRA COCHAIN TWISTS

We begin with some well known algebraic constructions, see for example the text [M2]. We will only need here the classical case $H = U(\mathcal{L})$ where $\mathcal{L}$ is a Lie algebra and $A = C^\infty(M)$ where M is a manifold, as the basis of the quantisation method. In that sense we use quantum group methods but the reader does not really need to know quantum group theory in any detail. This approach to quantisation as been recently used in [AM2, M3] for except that in the present paper the quantisation remains associative.

Given a Hopf algebra (H, S, Δ, ϵ) and an invertible $F \in H \otimes H$ with $(\epsilon \otimes \text{id})F = (\text{id} \otimes \epsilon)F = 1$, we can define a quasi-Hopf algebra $H_F = (H, \phi, S_F, \Delta_F, \epsilon, \alpha_F, \beta_F)$, with the same algebra and counit as H , by [D2]

$$(1) \quad \begin{aligned} \Delta_F h &= F \Delta h F^{-1}, \quad \phi = (1 \otimes F) \cdot (\text{id} \otimes \Delta) F \cdot (\Delta \otimes \text{id}) F^{-1} \cdot (F \otimes 1)^{-1}, \\ S_F &= S, \quad \alpha_F = (S F^{-(1)}) \cdot F^{-(2)}, \quad \beta_F = F^{(1)} S F^{(2)}. \end{aligned}$$

In addition if there is a quasitriangular structure $\mathcal{R}$ for H , then $\mathcal{R}_F = F_{21} \mathcal{R} F^{-1}$ for H_F . We will call such an $F \in H \otimes H$ a 2-cochain in general, and a 2-cocycle if $\phi = 1 \otimes 1 \otimes 1$.

The significance of the twisting construction is [M1] that it corresponds to an equivalence of categories. Thus, the category ${}_{H_F}\mathcal{M}$ of left modules over H_F is a monoidal category with tensor product operation $\otimes^F$ is defined using Δ_F . If $\phi = \phi^{(1)} \otimes \phi^{(2)} \otimes \phi^{(3)}$, the associator in the category is $\Phi_{VWZ}((v \otimes w) \otimes z) = \phi^{(1)} \triangleright v \otimes (\phi^{(2)} \triangleright w \otimes \phi^{(3)} \triangleright z)$. On the other hand, this category is equivalent to the category ${}_H\mathcal{M}$ of left modules over H via the functor $\mathcal{T} : {}_H\mathcal{M} \rightarrow {}_{H_F}\mathcal{M}$ which is just the identity on left H modules and on morphism. A monoidal functor also comes by definition with a natural transformation $\vartheta : \mathcal{T}V \otimes^F \mathcal{T}W \rightarrow \mathcal{T}(V \otimes W)$, given here by $\vartheta(\mathcal{T}(v) \otimes^F \mathcal{T}(w)) = \mathcal{T}(F^{-(1)} \triangleright v \otimes F^{-(2)} \triangleright w)$ [M1]. In this way, twisting the Hopf algebra by F deforms the entire category of modules and as such deforms any and all constructions in the category. This is the systematic 'twisting approach' to deformation quantisation that we use.

In particular, consider an algebra $A \in {}_H\mathcal{M}$. This includes the requirement that multiplication $: A \otimes A \rightarrow A$ is a morphism in the category, i.e the product is H -covariant (or A is an H -module algebra). Applying the above functor $\mathcal{T}$ immediately deforms the algebra to the same vector space $A_F = A$ and the product as a map $\mathcal{T}(A \otimes A) \rightarrow \mathcal{T}A$. Using the above natural transformation this implies a deformed product map making an algebra $A_F \in {}_{H_F}\mathcal{M}$ with multiplication $a \bullet b = (F^{-(1)} \triangleright a)(F^{-(2)} \triangleright b)$, and this is associative in the category as the image of the associativity law in the undeformed category. This module algebra cochain quantisation method was introduced in [DGM] and related papers at the time. Examples in the cocycle case also abound, e.g. [MO], but the cocycle case is not what is of interest in the present paper since in this case the associator ϕ is trivial. Neither case of 'module algebra twist' should be confused with Drinfeld's twist H_F of the Hopf algebra H itself.

One may go further and consider also the category $\mathcal{A} \subset {}_H\mathcal{M}$ of A -bimodules, which also have H -actions so that the multiplications $A \otimes V \rightarrow V$ and $V \otimes A \rightarrow V$ preserve the H -action for all $V \in {}_A\mathcal{M}$, and so on. Here we deform the multiplications by $a \bullet v = (F^{-(1)} \triangleright a).(F^{-(2)} \triangleright v)$ and $v \bullet a = (F^{-(1)} \triangleright v).(F^{-(2)} \triangleright a)$ for all $v \in V$ and $a \in A$. Similarly, if $\Omega(A)$ is an H -covariant differential calculus in the sense of noncommutative geometry (so there is for example an exterior derivative $d : A \rightarrow \Omega^1(A)$ where the latter is an A -bimodule and d obeys the Leibniz rule, etc. and all maps are morphisms in ${}_H\mathcal{M}$) then twisting any products by the action of F^{-1} gives a calculus $\Omega(A_F)$ covariant under H_F . This was used for example in [AM2].

To this existing theory we now add some first remarks needed for the semiclassical analysis. As mentioned, the above should be understood as extended over formal power-series in a parameter $\hbar$ or one may continue more algebraically (using a comodule twist version of the theory). Either way, we suppose that F^{-1} is expanded as a series

$$(2) \quad F^{-1} = 1 \otimes 1 + \hbar G^{(1)} + \hbar^2 G^{(2)} + O(\hbar^3) .$$

This can be inverted to give

$$(3) \quad F = 1 \otimes 1 - \hbar G^{(1)} + \hbar^2 ((G^{(1)})^2 - G^{(2)}) + O(\hbar^3) .$$

We can then compute

$$\begin{aligned} \phi &= (1 \otimes F).(\text{id} \otimes \Delta)F.(\Delta \otimes \text{id})F^{-1}.(F \otimes 1)^{-1} \\ &= 1 \otimes 1 \otimes 1 + \hbar [(\Delta \otimes \text{id})G^{(1)} + G^{(1)} \otimes 1 - 1 \otimes G^{(1)} - (\text{id} \otimes \Delta)G^{(1)}] \\ &\quad + \hbar^2 [(\Delta \otimes \text{id})G^{(2)} + G^{(2)} \otimes 1 + 1 \otimes ((G^{(1)})^2 - G^{(2)}) + (\text{id} \otimes \Delta)((G^{(1)})^2 - G^{(2)}) \\ &\quad + (\Delta \otimes \text{id})G^{(1)}.((G^{(1)} \otimes 1) - (\text{id} \otimes \Delta)G^{(1)}).(G^{(1)} \otimes 1) - (\text{id} \otimes \Delta)G^{(1)}.(\Delta \otimes \text{id})G^{(1)} \\ &\quad - (1 \otimes G^{(1)}).(G^{(1)} \otimes 1) - (1 \otimes G^{(1)}).(\Delta \otimes \text{id})G^{(1)} + (1 \otimes G^{(1)}).(\text{id} \otimes \Delta)G^{(1)}] + O(\hbar^3). \end{aligned}$$

Now consider a special case, where $G^{(1)} = \sum X \otimes Y$ (we will later suppress the summation sign), $\Delta X = 1 \otimes X + X \otimes 1$ and $\Delta Y = 1 \otimes Y + Y \otimes 1$. Then the order $\hbar$ part of ϕ is

$$1 \otimes X \otimes Y + X \otimes 1 \otimes Y + X \otimes Y \otimes 1 - 1 \otimes X \otimes Y - X \otimes Y \otimes 1 - X \otimes 1 \otimes Y = 0 .$$

The contribution from $G^{(1)}$ to the order $\hbar^2$ part of ϕ , using tildes to distinguish different copies of $G^{(1)}$, is

$$\begin{aligned}
& 1 \otimes X\tilde{X} \otimes Y\tilde{Y} + X\tilde{X} \otimes Y\tilde{Y} \otimes 1 + X\tilde{X} \otimes 1 \otimes Y\tilde{Y} + X\tilde{X} \otimes Y \otimes \tilde{Y} + X\tilde{X} \otimes \tilde{Y} \otimes Y \\
& + X\tilde{X} \otimes \tilde{Y} \otimes Y + \tilde{X} \otimes X\tilde{Y} \otimes Y - X\tilde{X} \otimes \tilde{Y} \otimes Y - X\tilde{X} \otimes Y\tilde{Y} \otimes 1 \\
& - X\tilde{X} \otimes Y \otimes \tilde{Y} - X\tilde{X} \otimes 1 \otimes Y\tilde{Y} - X \otimes Y\tilde{X} \otimes \tilde{Y} - X \otimes \tilde{X} \otimes Y\tilde{Y} - \tilde{X} \otimes X\tilde{Y} \otimes Y \\
& - \tilde{X} \otimes X \otimes Y\tilde{Y} - 1 \otimes X\tilde{X} \otimes Y\tilde{Y} + \tilde{X} \otimes X\tilde{Y} \otimes Y + \tilde{X} \otimes X \otimes Y\tilde{Y} \\
& = X\tilde{X} \otimes \tilde{Y} \otimes Y + \tilde{X} \otimes X\tilde{Y} \otimes Y - X \otimes Y\tilde{X} \otimes \tilde{Y} - X \otimes \tilde{X} \otimes Y\tilde{Y} \\
& = X\tilde{X} \otimes \tilde{Y} \otimes Y + \tilde{X} \otimes X\tilde{Y} \otimes Y - \tilde{X} \otimes \tilde{Y}X \otimes Y - X \otimes \tilde{X} \otimes Y\tilde{Y} \\
& = X\tilde{X} \otimes \tilde{Y} \otimes Y + \tilde{X} \otimes [X, \tilde{Y}] \otimes Y - X \otimes \tilde{X} \otimes Y\tilde{Y} .
\end{aligned}$$

To summarise, if we put, for $P \in H \otimes H$,

$$\partial P = 1 \otimes P - (\Delta \otimes \text{id})P + (\text{id} \otimes \Delta)P - P \otimes 1 .$$

then the expression for ϕ becomes

$$\phi = 1 \otimes 1 \otimes 1 + \hbar^2 (X\tilde{X} \otimes \tilde{Y} \otimes Y + \tilde{X} \otimes [X, \tilde{Y}] \otimes Y - X \otimes \tilde{X} \otimes Y\tilde{Y} - \partial G^{(2)}) + O(\hbar^3) .$$

Next note that, if $A, B, C, D \in H$ are also primitive (have linear coproducts like X, Y) then

$$\partial(AB \otimes CD) = AB \otimes C \otimes D + AB \otimes D \otimes C - A \otimes B \otimes CD - B \otimes A \otimes CD .$$

Hence if we were to put $G^{(2)} = (G^{(1)})^2/2$, then

$$(4) \quad \phi = 1 \otimes 1 \otimes 1 + \hbar^2 ([X, \tilde{X}] \otimes \tilde{Y} \otimes Y + 2\tilde{X} \otimes [X, \tilde{Y}] \otimes Y - X \otimes \tilde{X} \otimes [Y, \tilde{Y}]) / 2 + O(\hbar^3) .$$

The set of primitive elements in H form a Lie algebra in H which we can view as a Lie bialgebra with zero Lie cobracket. After twisting these elements acquire a Lie cobracket

$$\delta Z = \sum [X, Z] \otimes Y + X \otimes [Z, Y] - [Y, Z] \otimes X - Y \otimes [X, Z]$$

for all Z in the Lie algebra and together with the $\hbar^2$ part of ϕ form a quasi-Lie bialgebra. This is the infinitesimal object associated to the quasi-quantum group H_F .

3. POISSON-COMPATIBLE CONNECTIONS FROM COCHAIN TWISTS AT THE SEMICLASSICAL LEVEL

We start by briefly recalling the main ideas of [BM1]. As is well known, if one considers the quantisation of the functions $C^\infty(M)$ of a classical manifold M , the initial data usually specified is a Poisson structure defined by a bivector ω (in the symplectic case this is invertible with inverse (also denoted ω) a closed 2-form). Any flat deformation-quantisation $A_\hbar$ corresponds on looking at the leading part of its commutator

$$(5) \quad a \bullet b - b \bullet a = \hbar \{a, b\} + O(\hbar^2)$$

to a Poisson bracket $\{a, b\} = \omega(da, db)$. More recently, we considered the same question for a noncommutative differential calculus $\Omega(A_\hbar)$ quantizing the usual exterior algebra $\Omega(M)$ but in a slightly weaker than usual setting (without assuming associativity of products involving differential forms). We found that the initial data for this at least in the symplectic case was a compatible connection ∇ defined by

$$(6) \quad a \bullet db - db \bullet a = \hbar \nabla_{\hat{a}} db + O(\hbar^2).$$

Here $\hat{a}$ denotes the Hamiltonian vector field $\omega(da, \cdot) = \{a, \cdot\}$. Here ∇ is not necessarily well-defined in the Poisson case even along Hamiltonian vector fields; it could be called a partial connection where defined or we should speak more precisely of a 'preconnection' $\hat{\nabla}$ defined almost identically by

$$(7) \quad a \bullet db - db \bullet a = \hbar \hat{\nabla}_a db + O(\hbar^2)$$

(here $\hat{\nabla}$ was called γ in [BM1]). (In fact there is a more general notion of ‘contravariant connection’ which can also be used here, see[H]). The (Poisson)-compatibility condition is

$$(8) \quad \nabla_{\hat{a}} db - \nabla_{\hat{b}} da = d\{a, b\}$$

and under some mild conditions in the symplectic case becomes [BM1] that ∇ is a torsion free symplectic connection in the usual sense. Finally, the curvature and torsion of the connection

$$(9) \quad R_{\nabla}(\hat{a}, \hat{b}) = [\nabla_{\hat{a}}, \nabla_{\hat{b}}] - \nabla_{[\hat{a}, \hat{b}]}, \quad T_{\nabla}(\hat{a}, \hat{b}) = \nabla_{\hat{a}} \hat{b} - \nabla_{\hat{b}} \hat{a} - [\hat{a}, \hat{b}]$$

are defined in the usual way as for any connection. In terms of a preconnection the equations are more precisely

$$(10) \quad \hat{\nabla}_a db - \hat{\nabla}_b da = d\{a, b\}, \quad R_{\hat{\nabla}}(a, b) = [\hat{\nabla}_a, \hat{\nabla}_b] - \hat{\nabla}_{\{a, b\}}, \quad T_{\hat{\nabla}}(a, b) = \hat{\nabla}_a \hat{b} - \hat{\nabla}_b \hat{a} - [\hat{a}, \hat{b}]$$

where the last term in the curvature is in view of $[\hat{a}, \hat{b}] = \widehat{\{a, b\}}$. It was shown in [BM1] that the curvature coincides with the Jacobiator or obstruction to associativity for the differential calculus $\Omega(A_{\hbar})$ at the relevant lowest order. This was also observed recently in [H] where it was shown that the associative case corresponds to a zero-curvature (contravariant) connection, although we were not aware of this at the time of [BM1].

From a geometrical point of view, however, it would seem at the semiclassical level quite reasonable to consider symplectic or Poisson manifolds equipped with connections with curvature, which in quantisation terms would mean by [BM1] associative quantum algebras with nonassociative differential calculi. We have seen in Section 2 a general method to construct examples of such hybrid situations by means of cochain twists. Now we see what that amounts to at the semiclassical level.

3.1. Inducing connections by twisting. We consider the diffeomorphism group acting on the functions on a manifold M . This action extends to the vector fields and forms on the manifold, and infinitesimally the action is called the Lie derivative. A vector field X (i.e. an element of the Lie algebra of the diffeomorphism group) acts on forms by $\mathcal{L}_X \xi = \varpi_X(d\xi) + d(\varpi_X \xi)$, where ϖ_X is the interior product. If $\xi = \xi_{k_1 \dots k_n} dx^{k_1} \wedge \dots \wedge dx^{k_n}$, then

$$\begin{aligned} \mathcal{L}_X \xi &= X^p \xi_{k_1 \dots k_n, p} dx^{k_1} \wedge \dots \wedge dx^{k_n} + \xi_{k_1 \dots k_n} d(\varpi_X(dx^{k_1} \wedge \dots \wedge dx^{k_n})) \\ &= \left(X^j \xi_{k_1 \dots k_n, j} + X^j_{, k_i} \xi_{k_1 \dots k_{i-1}, j, k_{i+1} \dots k_n} \right) dx^{k_1} \wedge \dots \wedge dx^{k_n} \end{aligned}$$

We can therefore apply the ideas of Section 2 with $A = C^\infty(M)$ and $H = U(\text{diff}(M))$ acting via the Lie derivative on all tensorial objects. We will proceed with F, F^{-1} power-series having values in $U(\text{diff}(M)) \otimes U(\text{diff}(M))$, however we are interested in this section only in the differential geometry resulting from the semiclassical part and not in the formal construction of these objects.

We assume an expansion of the 2-cochain $F^{-1} = \text{id}^2 + \hbar G^{(1)} + O(\hbar^2)$, where $G^{(1)} = \sum X \otimes Y \in \text{diff}(M) \otimes \text{diff}(M)$, acts on $\Omega^n(M) \otimes \Omega^m(M)$ by $\xi \otimes \eta \mapsto \xi \otimes \eta + \hbar \sum \mathcal{L}_X \xi \otimes \mathcal{L}_Y \eta + O(\hbar^2)$. Then the commutator of two functions $a, b \in C^\infty(M)$ is

$$\begin{aligned} \{a, b\} &= \hbar \sum \left((\mathcal{L}_X a)(\mathcal{L}_Y b) - (\mathcal{L}_Y a)(\mathcal{L}_X b) \right) + O(\hbar^2) \\ (11) \quad &= \hbar \sum (X^i Y^j - X^j Y^i) a_{,i} b_{,j} + O(\hbar^2). \end{aligned}$$

Now we set $\omega = \sum (X \otimes_M Y - Y \otimes_M X) \in \text{diff}(M) \otimes_M \text{diff}(M)$ (putting summation subscripts on X and Y would only be confusing). This is antisymmetric, so we have $\omega \in \text{diff}(M) \wedge \text{diff}(M)$. We shall assume for convenience that we are in the symplectic case, where ω is invertible, and its inverse is a closed 2-form, otherwise more generally we assume that ω is a Poisson bivector, i.e. induces a Poisson bracket. A sufficient condition for the latter is that $\phi = 1 \otimes 1 \otimes 1$ when projected from $\otimes$ to $\otimes_M$. Another sufficient condition is that $G^{(1)}$ obeys the Classical Yang-Baxter equations, but neither is necessary and we do not assume them.

Now apply the same F^{-1} to deform the products of functions and 1-forms as explained in Section 2. This implies a connection ∇ resulting from the commutator of a function and a 1-form:

$$\begin{aligned}
[a, \xi_i dx^i] &= \hbar \sum X^k a_{,k} (\xi_{i,j} Y^j dx^i + \xi_i Y_{,j}^i dx^j) - (X \leftrightarrow Y) + O(\hbar^2) \\
&= \hbar \sum (X^k Y^j - X^j Y^k) a_{,k} \xi_{i,j} dx^i + \sum (X^k Y_{,j}^i - Y^k X_{,j}^i) \xi_i a_{,k} dx^j + O(\hbar^2) \\
&= \hbar \omega^{kj} a_{,k} \xi_{i,j} dx^i + \sum (X^k Y_{,j}^i - Y^k X_{,j}^i) \xi_i a_{,k} dx^j + O(\hbar^2) \\
&= \hbar \omega^{kj} a_{,k} \left(\xi_{i,j} dx^i + \sum \omega_{js} (X^s Y_{,p}^i - Y^s X_{,p}^i) \xi_i dx^p \right) + O(\hbar^2) \\
&= \hbar \omega^{kj} a_{,k} \left(\xi_{i,j} dx^i - \Gamma_{jp}^i \xi_i dx^p \right) + O(\hbar^2) .
\end{aligned}$$

Then the Christoffel symbols of the connection can be seen to be

$$(12) \quad \Gamma_{jp}^i = - \sum \omega_{js} (X^s Y_{,p}^i - Y^s X_{,p}^i) .$$

Proposition 3.1.1. *The connection is characterised by the equation, for $a \in C^\infty(M)$ and $\xi \in \Omega^1 M$:*

$$\nabla_a \xi = \sum X(a) (d\langle Y, \xi \rangle + \varpi_Y d\xi) - Y(a) (d\langle X, \xi \rangle + \varpi_X d\xi) .$$

Proof: Just substitute from (12) for the Christoffel symbols. $\heartsuit$

The condition that ω is closed is

$$\omega_{ir} \omega^{rs}{}_{,k} \omega_{sj} dx^k \wedge dx^i \wedge dx^j = 0 .$$

From [BM1] the connection is necessarily comapitable with the differential structure in the form

$$(13) \quad \frac{\partial \omega^{ij}}{\partial x^p} = \omega^{jq} \Gamma_{qp}^i - \omega^{iq} \Gamma_{qp}^j .$$

For the Christoffel symbols in (12) we have

$$\begin{aligned}
\omega^{jq} \Gamma_{qp}^i - \omega^{iq} \Gamma_{qp}^j &= \sum (X^i Y_{,p}^j - Y^i X_{,p}^j) - \sum (X^j Y_{,p}^i - Y^j X_{,p}^i) \\
&= \sum (X^i Y^j - X^j Y^i)_{,p} = \omega^{ij}{}_{,p} ,
\end{aligned}$$

as expected.

In summary, any manifold may potentially be quantised by choosing a cochain F with values in $U(\text{diff} M) \otimes U(\text{diff} M)[[\hbar]]$. The leading order of F which we have denoted $\sum X \otimes Y$ will induce a bivector which will not necessarily be a Poisson bivector (the quantised algebra may not necessarily be associative). However, if it is, we will also have induced a Poisson-compatible connection (or more precisely a preconnection). Conversely, given, say, a symplectic manifold M equipped with (as in Fedosov theory) a symplectic connection ∇ we can look for a suitable F inducing these initial data to lowest order and such that $\phi = 1 \otimes 1 \otimes 1$ when tensorised over $\otimes_M$. This provides an alternative and more categorical approach to the quantisation problem in the spirit of Fedosov theory but now having the merit of also quantising differential calculi and all other covariant constructions, albeit with potential nonassociativity.

3.2. The inverse problem for $\mathbb{R}^{2n}$. The ‘inverse problem’ of finding F even to lowest order (i.e. $\sum X \otimes Y$) such that a given Poisson bi-vector ω is obtained and a given symplectic or Poisson-compatible (pre)connection is obtained as above appears to be a tricky one. Here we will look at what is involved in the simplest possible case of $\mathbb{R}^{2n}$. We take the standard symplectic structure and note that torsion free symplectic connections for it are in 1-1 correspondence with totally symmetric Christoffel symbols $\Gamma_{abc} = \omega_{ad} \Gamma_{bc}^d$ [GRS].

We can easily make the canonical symplectic form for $\mathbb{R}^{2n}$ by adding $\hbar X \otimes Y$ terms to F^{-1} where X, Y are constant vector fields. By (12) these will give zero Christoffel symbols. But then

we can add further terms to F^{-1} of the form $\sum X \otimes Y = \sum f.U \otimes V - U \otimes f.V$, where f is a real valued function and U, V are constant vectors. Then

$$\Gamma_{ijp} = - \sum \omega_{ik} \omega_{js} f_{,p} (-U^s V^k - V^s U^k) .$$

If we set f to be the linear function $\omega_{pq} W^q x^p$, then

$$\Gamma_{ijp} = \omega_{ik} \omega_{js} \omega_{pq} \sum W^q (U^s V^k + V^s U^k) .$$

By adding terms of this form we can recreate any symplectic connection with constant Christoffel symbols by this form of F . Then the curvature is given by

$$\begin{aligned} R^a_{bcd} &= \omega^{me} \omega^{ag} (\Gamma_{edb} \Gamma_{gcm} - \Gamma_{ecb} \Gamma_{gdm}) \\ &= \omega^{me} \omega^{ag} \omega_{ek} \omega_{bq} \omega_{gp} \omega_{mr} \left(\omega_{ds} \omega_{cv} \sum W^q (U^s V^k + V^s U^k) \tilde{W}^r (\tilde{U}^v \tilde{V}^p + \tilde{V}^v \tilde{U}^p) \right. \\ &\quad \left. - \omega_{cs} \omega_{dv} \sum W^q (U^s V^k + V^s U^k) \tilde{W}^r (\tilde{U}^v \tilde{V}^p + \tilde{V}^v \tilde{U}^p) \right) \\ &= \omega_{bq} \omega_{kr} (\omega_{ds} \omega_{cv} - \omega_{cs} \omega_{dv}) \sum W^q (U^s V^k + V^s U^k) \tilde{W}^r (\tilde{U}^v \tilde{V}^a + \tilde{V}^v \tilde{U}^a) \\ &= \omega_{bq} \omega_{kr} (\omega_{ds} \omega_{cv} - \omega_{cs} \omega_{dv}) \sum W^q (U^s V^k \tilde{V}^v \tilde{U}^a + V^s U^k \tilde{U}^v \tilde{V}^a) \tilde{W}^r , \end{aligned}$$

where the tildes denote a second set of triples (U, V, W) and the sum is over both sets.

4. QUANTISING S^2 BY COCHAIN TWIST

Here we describe a simple example of the cochain quantisation method in Section 3. The covariance used for the twisting will be the Lorentz group and its action on S^2 , a nonlinear one related to spacetime physics (the sphere at infinity in Minkowski space). This induces a quantisation not related as far as we know to the representation theoretic coadjoint orbit examples given later.

4.1. Some nice vector fields on the 2-sphere. Our goal is to show how a natural covariance, cochain and hence quantisation arise in a nice way from the geometry of $S^2 = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = 1\}$. We use the standard inner product on $\mathbb{R}^3$.

Thus, given $\underline{v} \in \mathbb{R}^3$, we have the natural vector field $X[\underline{v}]$ defined by $X[\underline{v}](\underline{r}) = \underline{v} - \underline{r} \langle \underline{v}, \underline{r} \rangle$ (for $\underline{r} \in S^2$) which tangent to the sphere at $\underline{r}$. Also at each such point we have the orbital angular moment vector field $Y[\underline{v}](\underline{r}) = \underline{v} \times \underline{r}$, where $\times$ is the vector cross product. These vector fields are clearly well behaved under rotation; consider an orthogonal transformation $T \in O_3(\mathbb{R})$. Then $T(X[\underline{v}])(\underline{r})$ is by definition

$$T(X[\underline{v}](T^{-1}\underline{r})) = T\underline{v} - T(T^{-1}\underline{r}) \langle \underline{v}, T^{-1}\underline{r} \rangle = T\underline{v} - \underline{r} \langle T\underline{v}, \underline{r} \rangle = X[T\underline{v}](\underline{r}) .$$

Also we have $T(Y[\underline{v}])(\underline{r})$ equal to

$$T(Y[\underline{v}](T^{-1}\underline{r})) = T(\underline{v} \times T^{-1}\underline{r}) = \det(T).T\underline{v} \times \underline{r} = \det(T).Y[T\underline{v}](\underline{r}) ,$$

where the determinant enters by the change in sign of the vector product under a change in orientation of $\mathbb{R}^3$.

The Lie bracket of two vector fields is defined as usual by $[X, Y]^i = Y^i_{,j} X^j - X^i_{,j} Y^j$. Thus

$$\begin{aligned} [Y[\underline{v}], Y[\underline{w}]](\underline{r}) &= \underline{w} \times (\underline{v} \times \underline{r}) - \underline{v} \times (\underline{w} \times \underline{r}) \\ &= \langle \underline{w}, \underline{r} \rangle \underline{v} - \langle \underline{w}, \underline{v} \rangle \underline{r} - \langle \underline{v}, \underline{r} \rangle \underline{w} + \langle \underline{v}, \underline{w} \rangle \underline{r} \\ &= \langle \underline{w}, \underline{r} \rangle \underline{v} - \langle \underline{v}, \underline{r} \rangle \underline{w} , \\ [Y[\underline{v}], X[\underline{w}]](\underline{r}) &= -\underline{r} \langle \underline{w}, Y[\underline{v}](\underline{r}) \rangle - Y[\underline{v}](\underline{r}) \langle \underline{w}, \underline{r} \rangle - \underline{v} \times X[\underline{w}](\underline{r}) \\ &= -\underline{r} \langle \underline{w}, \underline{v} \times \underline{r} \rangle - \underline{v} \times \underline{r} \langle \underline{w}, \underline{r} \rangle - \underline{v} \times (\underline{w} - \underline{r} \langle \underline{w}, \underline{r} \rangle) \\ &= -\underline{r} \langle \underline{w}, \underline{v} \times \underline{r} \rangle - \underline{v} \times \underline{w} = \underline{r} \langle \underline{v} \times \underline{w}, \underline{r} \rangle - \underline{v} \times \underline{w} \\ &= -X[\underline{v} \times \underline{w}](\underline{r}) , \end{aligned}$$

$$\begin{aligned}
[X[\underline{v}], X[\underline{w}]](\underline{r}) &= -\underline{r} \langle \underline{w}, X[\underline{v}](\underline{r}) \rangle - X[\underline{v}](\underline{r}) \langle \underline{w}, \underline{r} \rangle + \underline{r} \langle \underline{v}, X[\underline{w}](\underline{r}) \rangle + X[\underline{w}](\underline{r}) \langle \underline{v}, \underline{r} \rangle \\
&= -\underline{r} \langle \underline{w}, \underline{v} \rangle - \underline{v} \langle \underline{w}, \underline{r} \rangle + \underline{r} \langle \underline{v}, \underline{w} \rangle + \underline{w} \langle \underline{v}, \underline{r} \rangle \\
&\quad + \underline{r} \langle \underline{w}, \underline{r} \langle \underline{v}, \underline{r} \rangle \rangle + \underline{r} \langle \underline{v}, \underline{r} \rangle \langle \underline{w}, \underline{r} \rangle - \underline{r} \langle \underline{v}, \underline{r} \langle \underline{w}, \underline{r} \rangle \rangle - \underline{r} \langle \underline{w}, \underline{r} \rangle \langle \underline{v}, \underline{r} \rangle \\
&= -\underline{v} \langle \underline{w}, \underline{r} \rangle + \underline{w} \langle \underline{v}, \underline{r} \rangle
\end{aligned}$$

whereas

$$Y[\underline{v} \times \underline{w}](\underline{r}) = (\underline{v} \times \underline{w}) \times \underline{r} = \langle \underline{r}, \underline{v} \rangle \underline{w} - \langle \underline{r}, \underline{w} \rangle \underline{v} .$$

Hence we have

$$(14) Y[\underline{v}], Y[\underline{w}] = -Y[\underline{v} \times \underline{w}] , \quad [Y[\underline{v}], X[\underline{w}]] = -X[\underline{v} \times \underline{w}] , \quad [X[\underline{v}], X[\underline{w}]] = Y[\underline{v} \times \underline{w}] .$$

In other words, the Y fields generate rotations and the X generate boosts of the Lie algebra $\mathcal{L} = so(1, 3) \subset \text{diff}(S^2)$. This action has the physical interpretation mentioned above and will be used to induce the quantisation.

4.2. The first order rotation invariant 2-cochain. Set

$$\begin{aligned}
X_1 &= X[(1, 0, 0)] , \quad X_2 = X[(0, 1, 0)] , \quad X_3 = X[(0, 0, 1)] , \\
Y_1 &= Y[(1, 0, 0)] , \quad Y_2 = Y[(0, 1, 0)] , \quad Y_3 = Y[(0, 0, 1)] .
\end{aligned}$$

Using the matrix coefficients of $T \in O_3(\mathbb{R})$ in the standard basis,

$$T(X_i) = \sum_j T_{ji} X_j , \quad T(Y_i) = \det T \cdot \sum_j T_{ji} Y_j ,$$

so under a rotation each X_i is sent to a linear combination of the X_j with (the important bit) constant coefficients, not general functions on S^2 , and likewise with the Y_i . Using these notations, for any 3×3 real matrix κ , emphasising the fact that the tensor product is over $\mathbb{R}$, we take the lowest order part of F, F^{-1} in the form

$$G_{\kappa}^{(1)} = \kappa_{ij} X_i \otimes_{\mathbb{R}} Y_j .$$

Then,

$$(T \otimes T)G_{\kappa}^{(1)} = \det T \cdot \kappa_{ij} T_{ki} X_k \otimes_{\mathbb{R}} T_{sj} Y_s = \det T \cdot T_{ki} \kappa_{ij} T_{js}^T X_k \otimes_{\mathbb{R}} Y_s = \det T \cdot G_{T\kappa T^{-1}}^{(1)}$$

Hence if we want $\Upsilon[\kappa]$ to be rotation invariant, then we would like $T\kappa T^{-1} = \kappa$ for all $T \in SO_3(\mathbb{R})$; we therefore take κ to be (half) the identity matrix. Of course it will still have its sign changed by orientation reversing $T \in O_3(\mathbb{R})$, but this is also true of a rotation invariant symplectic form on S^2 , so that this is exactly what we want. This last consideration also excludes terms of the form $X \otimes X$ and $Y \otimes Y$ in our ansatz for $G^{(1)}$.

We are therefore led by rotational considerations to $G^{(1)} = \frac{1}{2} X_i \otimes Y_i$, which we use henceforth. We use (11) to find the corresponding Poisson structure. As it is rotation invariant, we only have to evaluate $G^{(1)}$ at the point $(0, 0, 1)$:

$$\begin{aligned}
2G^{(1)}(0, 0, 1) &= X_1(0, 0, 1) \otimes Y_1(0, 0, 1) + X_2(0, 0, 1) \otimes Y_2(0, 0, 1) \\
&= -(1, 0, 0) \otimes (0, 1, 0) + (0, 1, 0) \otimes (1, 0, 0) .
\end{aligned}$$

This means that, were we to reduce to $\otimes_{C(S^2)}$, we would get the Poisson structure corresponding to the usual symplectic form on S^2 .

4.3. The connection. It will be convenient to choose coordinates $(x, y) \in \mathbb{R}^2$ for the hemisphere $\{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = 1 \text{ and } z > 0\}$. Then the component vector fields are given by

$$\begin{aligned} X_1(x, y) &= (1 - x^2, -xy), & X_2(x, y) &= (-xy, 1 - y^2), & X_3(x, y) &= (-xz, -yz), \\ Y_1(x, y) &= (0, -z), & Y_2(x, y) &= (z, 0), & Y_3(x, y) &= (-y, x). \end{aligned}$$

The Poisson tensor corresponding to the unique $G^{(1)}$ found above is

$$2\omega = \sum_i \left(X_i \otimes_{C(S^2)} Y_i - Y_i \otimes_{C(S^2)} X_i \right),$$

so, numbering the coordinates $x^1 = x$ and $x^2 = y$, ω^{ij} is antisymmetric and $\omega^{12}(x, y) = -z$. Taking the inverse matrix gives $\omega_{12} = 1/z$. From (12) the Christoffel symbols are (with summation sign suppressed)

$$\begin{aligned} 2\Gamma_{1p}^i &= \omega_{1s} (Y_k^s X_{k,p}^i - X_k^s Y_{k,p}^i) = (Y_k^2 X_{k,p}^i - X_k^2 Y_{k,p}^i)/z, \\ 2\Gamma_{2p}^i &= \omega_{2s} (Y_k^s X_{k,p}^i - X_k^s Y_{k,p}^i) = -(Y_k^1 X_{k,p}^i - X_k^1 Y_{k,p}^i)/z. \end{aligned}$$

In more detail,

$$\begin{aligned} 2\Gamma_{1p}^1 &= (Y_k^2 X_{k,p}^1 - X_k^2 Y_{k,p}^1)/z \\ &= (-z(1 - x^2)_{,p} + x(-xz)_{,p} - (1 - y^2)z_{,p} - yzy_{,p})/z, \\ 2\Gamma_{1p}^2 &= (Y_k^2 X_{k,p}^2 - X_k^2 Y_{k,p}^2)/z \\ &= (-z(-xy)_{,p} + x(-yz)_{,p} + xy(-z)_{,p} + yzx_{,p})/z, \\ 2\Gamma_{2p}^1 &= (X_k^1 Y_{k,p}^1 - Y_k^1 X_{k,p}^1)/z \\ &= (-xy z_{,p} + xz y_{,p} - z(-xy)_{,p} + y(-xz)_{,p})/z, \\ 2\Gamma_{2p}^2 &= (X_k^1 Y_{k,p}^2 - Y_k^1 X_{k,p}^2)/z \\ &= ((1 - x^2)(-z)_{,p} - xz x_{,p} - z(1 - y^2)_{,p} + y(-yz)_{,p})/z. \end{aligned}$$

Then we get, using $z_{,1} = -x/z$ and $z_{,2} = -y/z$,

$$\begin{aligned} 2\Gamma_{1p}^1 &= (zx x_{,p} - x^2 z_{,p} - (1 - y^2)z_{,p} - yzy_{,p})/z, \\ 2\Gamma_{11}^1 &= (zx + x^3/z + (1 - y^2)x/z)/z = x(z^2 + x^2 + 1 - y^2)/z^2, \\ \Gamma_{11}^1 &= x(1 - y^2)/z^2, \\ 2\Gamma_{12}^1 &= (x^2 y/z + (1 - y^2)y/z - yz)/z = y(x^2 + 1 - y^2 - z^2)/z^2, \\ \Gamma_{12}^1 &= x^2 y/z^2, \\ 2\Gamma_{1p}^2 &= (z(xy)_{,p} - x(yz)_{,p} - xyz_{,p} + yzx_{,p})/z = 2(zx_p y - xyz_{,p})/z, \\ \Gamma_{11}^2 &= y(1 - y^2)/z^2, \\ \Gamma_{12}^2 &= xy^2/z^2, \\ 2\Gamma_{2p}^1 &= (-xyz_{,p} + xz y_{,p} + z(xy)_{,p} - y(xz)_{,p})/z = 2(xz y_{,p} - yxz_{,p})/z, \\ \Gamma_{21}^1 &= x^2 y/z^2, \\ \Gamma_{22}^1 &= x(1 - x^2)/z^2, \\ 2\Gamma_{2p}^2 &= (-(1 - x^2)z_{,p} - xz x_{,p} + zyy_{,p} - y^2 z_{,p})/z, \\ 2\Gamma_{22}^2 &= ((1 - x^2)y/z + zy + y^3/z)/z = y(1 - x^2 + z^2 + y^2)/z^2, \\ \Gamma_{22}^2 &= y(1 - x^2)/z^2, \\ 2\Gamma_{21}^2 &= ((1 - x^2)x/z - xz + y^2 x/z)/z = x(1 - x^2 - z^2 + y^2)/z^2, \\ \Gamma_{21}^2 &= xy^2/z^2. \end{aligned}$$

From this we see that the connection is torsion free, and since it is also compatible with the differential structure, by [BM1, Sec. 3] it is also symplectic.

4.4. Metric compatability. The metric induced from the standard embedding in $\mathbb{R}^3$ with the standard inner product is, in (x, y) coordinates,

$$g = \{g_{ij}\} = \frac{1}{z^2} \begin{pmatrix} 1-y^2 & xy \\ xy & 1-x^2 \end{pmatrix}.$$

If we organise the Christoffel symbols into the matrices

$$N_k = \begin{pmatrix} \Gamma_{k1}^1 & \Gamma_{k1}^2 \\ \Gamma_{k2}^1 & \Gamma_{k2}^2 \end{pmatrix},$$

then, using matrix multiplication,

$$\nabla_k g_{ij} = \frac{\partial g_{ij}}{\partial x^k} - N_k \cdot g_{ij} - g_{ij} \cdot N_k^T.$$

In our case

$$N_1 = \frac{1}{z^2} \begin{pmatrix} x(1-y^2) & y(1-y^2) \\ x^2 y & x y^2 \end{pmatrix}, \quad N_2 = \frac{1}{z^2} \begin{pmatrix} x^2 y & x y^2 \\ x(1-x^2) & y(1-x^2) \end{pmatrix},$$

and using this it can quickly be verified that the covariant derivatives of the metric vanish. Since the induced connection above was torsion free, it must be the usual Levi-Civita connection on S^2 .

4.5. The curvature. In a coordinate frame, the curvature is given by

$$R^l_{ijk} = \frac{\partial \Gamma_{ki}^l}{\partial x^j} - \frac{\partial \Gamma_{ji}^l}{\partial x^k} + \Gamma_{ki}^m \Gamma_{jm}^l - \Gamma_{ji}^m \Gamma_{km}^l.$$

At the point $x = y = 0$ we find that, to first order in x and y , $\Gamma_{11}^1 = \Gamma_{22}^1 = x$, $\Gamma_{11}^2 = \Gamma_{22}^2 = y$ and all other Christoffel symbols vanish. Then, at that point,

$$R^l_{ijk} = \frac{\partial \Gamma_{ki}^l}{\partial x^j} - \frac{\partial \Gamma_{ji}^l}{\partial x^k} = \delta_j^l \delta_{ki} - \delta_k^l \delta_{ji}.$$

Now we set $G^{(1)} = X_i \otimes Y_i / 2$ (summed over i) as above, and $G^{(2)} = (G^{(1)})^2 / 2$, then from (4)

$$\begin{aligned} \phi &= 1 \otimes 1 \otimes 1 + \hbar^2 ([X_i, X_j] \otimes Y_j \otimes Y_i + 2 X_j \otimes [X_i, Y_j] \otimes Y_i \\ &\quad - X_i \otimes X_j \otimes [Y_i, Y_j]) / 8 + O(\hbar^3) \\ &= 1 \otimes 1 \otimes 1 + \hbar^2 (Y[\underline{e}_i \times \underline{e}_j] \otimes Y_j \otimes Y_i - 2 X_j \otimes X[\underline{e}_i \times \underline{e}_j] \otimes Y_i \\ &\quad + X_i \otimes X_j \otimes Y[\underline{e}_i \times \underline{e}_j]) / 8 + O(\hbar^3), \end{aligned} \tag{15}$$

where the $\underline{e}_i$ are the usual basis vectors. Now we can expand the summations:

$$\begin{aligned} X_i \otimes X_j \otimes Y[\underline{e}_i \times \underline{e}_j] &= X_{[1]} \otimes X_{[2]} \otimes Y_{[3]} - X_{[1]} \otimes X_{[3]} \otimes Y_{[2]} + X_{[2]} \otimes X_{[3]} \otimes Y_{[1]} \\ &\quad - X_{[2]} \otimes X_{[1]} \otimes Y_{[3]} + X_{[3]} \otimes X_{[1]} \otimes Y_{[2]} - X_{[3]} \otimes X_{[2]} \otimes Y_{[1]}, \\ X_j \otimes X[\underline{e}_i \times \underline{e}_j] \otimes Y_i &= X_{[2]} \otimes X_{[3]} \otimes Y_{[1]} - X_{[1]} \otimes X_{[3]} \otimes Y_{[2]} - X_{[3]} \otimes X_{[2]} \otimes Y_{[1]} \\ &\quad + X_{[1]} \otimes X_{[2]} \otimes Y_{[3]} + X_{[3]} \otimes X_{[1]} \otimes Y_{[2]} - X_{[2]} \otimes X_{[1]} \otimes Y_{[3]}, \end{aligned}$$

and using this (15) simplifies to

$$\phi = 1 \otimes 1 \otimes 1 + \hbar^2 (Y[\underline{e}_i \times \underline{e}_j] \otimes Y_j - X_j \otimes X[\underline{e}_i \times \underline{e}_j]) \otimes Y_i / 8 + O(\hbar^3). \tag{16}$$

Again expanding the summations, and assigning a name to part of (16),

$$\begin{aligned} \psi &= (Y[\underline{e}_i \times \underline{e}_j] \otimes Y_j - X_j \otimes X[\underline{e}_i \times \underline{e}_j]) \otimes Y_i \\ &= (Y_{[3]} \otimes Y_{[2]} - Y_{[2]} \otimes Y_{[3]} - X_{[2]} \otimes X_{[3]} + X_{[3]} \otimes X_{[2]}) \otimes Y_{[1]} \\ &\quad + (Y_{[1]} \otimes Y_{[3]} - Y_{[3]} \otimes Y_{[1]} - X_{[3]} \otimes X_{[1]} + X_{[1]} \otimes X_{[3]}) \otimes Y_{[2]} \\ &\quad + (Y_{[2]} \otimes Y_{[1]} - Y_{[1]} \otimes Y_{[2]} - X_{[1]} \otimes X_{[2]} + X_{[2]} \otimes X_{[1]}) \otimes Y_{[3]} \end{aligned}$$

Lemma 4.5.1. *If we take π to be the reduction to $\otimes_{C^\infty(S^2)}$, then $\pi\psi = 0$.*

Proof. This is a rather long calculation done with Mathematica. Details are omitted. $\heartsuit$

This corresponds to the multiplication of functions being associative to $O(\hbar^2)$. This critical fact justifies our choice of $G^{(2)} = (G^{(1)})^2/2$. Note that in this case we can use (3) to calculate

$$F = 1 \otimes 1 - \hbar G^{(1)} + \hbar^2 (G^{(1)})^2/2 + O(\hbar^3)$$

Note that this would be consistent with $F^{-1} = e^{\hbar G^{(1)}} = e^{\frac{\hbar}{2} X_i \otimes Y_i}$ but also note that $so(1,3)$ is not Abelian such an exponential form will not have $\phi = 1 \otimes 1 \otimes 1$, and indeed this is not true even at order $\hbar^2$. However, we see that when projected over $C^\infty(S^2)$ we do have that ϕ is effectively trivial at this order on functions; in other words the example demonstrates the hybrid set up of our paper at this order.

4.6. The deformed algebra. Following (2), if $G^{(1)} = X_i \otimes Y_i/2$ (summation suppressed) and $G^{(2)} = (G^{(1)})^2/2$, then we have

$$f \bullet g = fg + \hbar (X_i \triangleright f)(Y_i \triangleright g)/4 + \hbar^2 (X_i X_j \triangleright f)(Y_i Y_j \triangleright g)/8 + O(\hbar^3).$$

It will be convenient to continue to use the coordinates in 4.3, in which case, using subscripts for partial differentiation,

$$\begin{aligned} (X_i \triangleright f)(Y_i \triangleright g) &= ((1 - x^2)f_x - xyf_y)(-zg_y) + (-xyf_x + (1 - y^2)f_y)zg_x \\ &\quad + (-xz f_x - yz f_y)(-yg_x + xg_y) \\ &= z(f_y g_x - f_x g_y). \end{aligned}$$

With rather more work, we get the following formula:

$$\begin{aligned} (x^a y^b) \bullet (x^c y^d) &= x^{a+c} y^{b+d} + \hbar z x^{a+c-1} y^{b+d-1} (bc - ad)/2 \\ &\quad + \hbar^2 x^{a+c-2} y^{b+d-2} \left((bc(b-1)(c-1) + ad(a-1)(d-1) - 2abcd) \right. \\ &\quad + y^2 (ac - a^2 c + b^2 c - a c^2 - b^2 c^2 + 2abcd + a d^2 - a^2 d^2) \\ &\quad + x^2 (b c^2 - b^2 c^2 + a^2 d + b d - b^2 d + 2abcd - a^2 d^2 - b d^2) \\ &\quad \left. + x^2 y^2 (a+b)(c+d)(1+a+b+c+d) \right) /8 + O(\hbar^3). \end{aligned}$$

The second order part of $f \bullet g$, evaluated at $x = y = 0$, is one eighth of

$$\frac{\partial^2 f}{\partial x^2} \frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 f}{\partial y^2} \frac{\partial^2 g}{\partial x^2} - 2 \frac{\partial^2 f}{\partial x \partial y} \frac{\partial^2 g}{\partial x \partial y} - \frac{\partial f}{\partial x} \frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \frac{\partial g}{\partial y}.$$

Using rotation invariance, and the fact that all the Christoffel symbols vanish at $x = y = 0$, we see that this is

$$\omega^{ij} \omega^{kl} (\nabla_i f_{,k}) (\nabla_j g_{,l}) - g^{ij} f_{,i} g_{,j}.$$

This quantisation can be compared the Fedosov one for S^2 with the above symplectic structure and symplectic connection. We see that the second order part is not that given by the Fedosov method, as that does not have the $g^{ij} f_{,i} g_{,j}$ term. Note that while Fedosov gives a prescription for a quantisation that is associative on the functions to all orders from the symplectic form and connection, this is not necessarily a unique quantisation. However we expect that our second order term $G^{(2)}$ may have to be modified to allow extension to all orders in $\hbar$ as an associative algebra, and it is not obvious that this could be done within our existing 6 dimensional subalgebra of the vector fields.

5. ENVELOPING ALGEBRAS $U(g)$ AS COCHAIN TWISTS

As an important application of the ideas in Section 3, we consider $M = \mathfrak{g}^*$, the dual of a Lie algebra, equipped with its standard Kirillov-Kostant Poisson structure $\{v, w\} = [v, w]$. Here $S(\mathfrak{g}) = \mathbb{C}[\mathfrak{g}^*]$ i.e. we work with polynomial functions as generated by $v \in \mathfrak{g}$ viewed as linear functions on $\mathfrak{g}^*$.

5.1. The cochain to lowest order. To express these canonical data as induced by a cochain twist, we seek suitable vector fields to define $G^{(1)} = \sum X \otimes Y$. Some natural vector fields are $\mathfrak{g}$ itself acting by ad as mentioned above, i.e. the vector fields for the coadjoint action on $\mathfrak{g}^*$ from a geometrical point of view. Then there is $\mathfrak{g}^*$ acting by interior product on $S(\mathfrak{g})$, which is to say usual differentiation on $\mathfrak{g}^*$. These classes of vector fields generated a sub-Lie algebra $\mathcal{L} = \mathfrak{g} \ltimes \mathfrak{g}^* \subset \text{diff}(\mathfrak{g}^*)$ that turns out to be sufficient to induce the desired quantisation. Thus we take $H = U(\mathfrak{g} \ltimes \mathfrak{g}^*) = U(\mathfrak{g}) \ltimes S(\mathfrak{g}^*)$ acting covariantly on $A = S(\mathfrak{g})$.

Choose a basis $\{e_i\}$ in $\mathfrak{g}$, and a dual basis $\{e^i\}$ in $\mathfrak{g}^*$, and set

$$F^{-1} = 1 \otimes 1 + \alpha \hbar e_i \otimes e^i + \beta \hbar e^i \otimes e_i + O(\hbar^2).$$

Then

$$\omega(dv, dw) = (\alpha - \beta) \sum e_i(v) e^i(w) - e^i(v) e_i(w) = -2(\alpha - \beta)[v, w]$$

so we obtain the Kirillov-Kostant bracket with $\beta - \alpha = \frac{1}{2}$. As a first consequence:

Proposition 5.1.1. $\mathfrak{g}^*$ also has on it a canonical Poisson-compatible preconnection

$$\hat{\nabla}_v dw = \frac{1}{2} d[v, w]$$

where $\hat{v} = \text{ad}_v$ is the adjoint action viewed as a vector field (in classical differential geometry, this is a derivation on $S(\mathfrak{g})$). The curvature and torsion are

$$R(v, w)dz = -\frac{1}{4}[[v, w], z], \quad \langle T(v, w), dz \rangle = \frac{1}{2}[[v, w], z]$$

Proof. This comes out of the construction by using F to deform the differential calculus, and the leading order part $\sum X \otimes Y$ found above. With hindsight one may check independently using the axioms in [BM1] that this is indeed a canonical Poisson-compatible preconnection for the Kirillov-Kostant bracket. We compute its curvature as

$$R(v, w)(dz) = (\hat{\nabla}_v \hat{\nabla}_w - \hat{\nabla}_w \hat{\nabla}_v - \nabla_{\widehat{\{v, w\}}})(dz) = d([v, [w, z]] - [w, [v, z]] - 2[[v, w], z])/4$$

with the result stated in view of the Jacobi identity in the Lie algebra. Similarly

$$\langle T(v, w), dz \rangle = \langle \hat{\nabla}_v \hat{w} - \hat{\nabla}_w \hat{v} - [\hat{v}, \hat{w}], dz \rangle = \hat{v}(\hat{w}(z)) - \langle \hat{w}, \hat{\nabla}_v dz \rangle - (v \leftrightarrow w) - [[v, w], z]$$

with the result stated again on using the Jacobi identity. Incidentally, leaving out the 1/2 gives a preconnection with zero curvature, but it is not Poisson-compatible. $\heartsuit$

Note that the most general translation-invariant $\hat{\nabla}$ in this quantisation is given by the analysis of [BM1] as of the form

$$\hat{\nabla}_v dw = \frac{1}{2} d[v, w] + d\hat{\Xi}(v, w)$$

where $\hat{\Xi} : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ is some symmetric linear map. This follows from regarding $\mathfrak{g}^*$ as an Abelian Lie group and applying the theory in [BM1, Sec. 4.1]. The operations L^* and R^* translating differentials back to the origin are trivial so that $\hat{\nabla}_v dw = d\Xi(v, w)$ is defined by a map Ξ with arbitrary symmetric part, which we denote $\hat{\Xi}$, and antisymmetric part the same as above. On the other hand, if we further demand background ‘rotational’ invariance in the sense of ad-invariance under

$\mathfrak{g}$ (which becomes covariance of the calculus under the quantum double $D(U(\hat{\mathfrak{g}}))$ after quantisation) this corresponds to $\hat{\Xi}$ symmetric and ad-invariant.

Theorem 5.1.2. *For all simple $\mathfrak{g}$ other than sl_n , $n > 2$ the canonical preconnection in Proposition 5.1.1 is the only translation and $\mathfrak{g}$ -invariant one on $S(\mathfrak{g}) = \mathbb{C}[\mathfrak{g}^*]$. For $\mathfrak{g} = sl_n$, $n > 2$ there is a 1-parameter moduli space of such covariant $\hat{\nabla}$ but they all have curvature. Hence for all simple $\mathfrak{g}$ any covariant differential calculus on $U_h(\mathfrak{g})$ with classical dimensions is necessarily nonassociative.*

Proof. By the same arguments from invariant theory as in the proof of [BM1, Theorem 4.20] (but now in a different context), basically from Kostant's work, there is no nonzero symmetric ad-invariant map $\hat{\Xi} : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ for $\mathfrak{g}$ simple other than for sl_n , $n > 2$. Hence $\hat{\Xi} = 0$ and $\hat{\nabla}$ has to be the one in Proposition 5.1.1.

For sl_n , $n > 2$ one has the possibility of a 1-parameter family via the invariant totally symmetric trilinear form viewed as the map $\hat{\Xi}$. In this case

$$R(v, w)dz = -\frac{1}{4}d[[v, w], z] + d\left(\hat{\Xi}(v, \hat{\Xi}(w, z)) - \hat{\Xi}(w, \hat{\Xi}(v, z))\right)$$

since the terms linear in $\hat{\Xi}$ cancel using its ad-invariance. We have to show that there are always v, w, z with the curvature expression nonzero. To do this, note that the symmetric trilinear is a cubic polynomial on sl_n which on $v \in sl_n$ has value $I(v, v, v) = I(v)$, say (e.g. for sl_3 we have $I(v) = \det(v)$). We can reconstruct the full trilinear from this by polarisation, e.g.

$$I(v, w, w) = \frac{1}{6}(I(v + 2w) - 2I(v + w) + I(v)) - I(w)$$

and we define $\hat{\Xi}(v, w) = I(v, w, e_i)e_j\kappa^{ij}$ where κ^{ij} is the inverse matrix of the Killing form (not necessarily normalised). We fix v, w diagonal (i.e. in the standard Cartan subalgebra of sl_n) and focus on

$$R(v, w)dw = I(v, e_i, w)I(w, v, e_j)\kappa^{ij} - I(w, e_i, w)I(v, v, e_j)\kappa^{ij}.$$

We will show that this can be arranged to be non-zero. Note that if t lies in the Cartan subalgebra and $z \in sl_n$, then ad-invariance $I([t, v], w, z) + I(v, [t, w], z) + I(v, w, [t, z]) = 0$ means $I(v, w, [t, z]) = 0$. We conclude that if z is a root vector then, $I(v, w, z) = 0$ (since $[t, z]$ is a nonzero multiple of z). Hence $I(v, w, z)$ vanishes for all z in the space spanned by the nonzero root vectors, which is to say the orthogonal complement of the Cartan with respect to the Killing form (it is the space of matrices in sl_n with zero diagonal). Hence we let $\{e_a\}$ be a basis of the Cartan subalgebra completed to a basis of sl_n taken from this complement. It means that we can compute $R(v, w)dw$ using only a sum over the e_a, e_b in place of e_i, e_j in the expression above. For sl_3 we take $t_1 = e_{11} - e_{22}$, $t_2 = e_{22} - e_{33}$ in the Cartan. Then $e_1 = t_1$ and $e_2 = t_1 + \frac{1}{2}t_2 = \frac{1}{2}(e_{11} + e_{22}) - e_{33}$ are a basis with $\kappa^{ab} = \text{diag}(1/2, 2/3)$. We also compute

$$I_{111} \equiv I(t_1, t_1, t_1) = I_{222} \equiv I(t_2, t_2, t_2) = -1, \quad I_{112} \equiv I(t_1, t_1, t_2) = \frac{3}{2}, \quad I_{122} \equiv I(t_1, t_2, t_2) = \frac{5}{6}$$

using the polarisation formula above. Hence setting $v = t_1$, $w = t_2$ we compute

$$\begin{aligned} R(v, w)dw &= \frac{1}{2}I(t_1, t_2, t_1)^2 + \frac{2}{3}I(t_1, t_2, t_1 + \frac{1}{2}t_2)^2 \\ &\quad - \frac{1}{2}I(t_1, t_1, t_1)I(t_2, t_2, t_1) - \frac{2}{3}I(t_1, t_1, t_1 + \frac{1}{2}t_2)I(t_2, t_2, t_1 + \frac{1}{2}t_2) \\ &= \frac{1}{2}I_{111}^2 + \frac{2}{3}(I_{112} + \frac{1}{2}I_{122})^2 - \frac{1}{2}I_{111}I_{122} - \frac{2}{3}(I_{111} + \frac{1}{2}I_{112})(I_{122} + \frac{1}{2}I_{222}) > 0 \end{aligned}$$

for the values stated. This proves the result for sl_3 . For sl_n the trilinear is given by $I(v) = \sum_{i < j < k} v^i v^j v^k$ in terms of the diagonal entries of v in the Cartan. We take the same $v = t_1$, $w = t_2$

as above but viewed in the standard way inside sl_n rather than sl_3 and the e_1, e_2 completed to a diagonal basis for κ . Then the computation reduces to the same one as above for sl_3 . $\heartsuit$

In summary, these results tells us that for simple $\mathfrak{g}$ we are going to necessarily have to work with nonassociative differentials, and for all Lie algebras $\mathfrak{g}$ the canonical ‘universal’ choice (which is often the only choice) at the lowest order level is the one in Proposition 5.1.1. We therefore focus on this and have seen that it is indeed given by a cochain twist at lowest order. We next want to extend Proposition 5.1.1 to find F, F^{-1} at least to order $O(\hbar^3)$. To do this we first look at the product of $U_\hbar(\mathfrak{g})$ on monomials. Computations have been done with MATHEMATICA.

5.2. The Campbell-Baker-Hausdorff product. We consider which F induce not only the above semiclassical data but the actual product of $U_\hbar(\mathfrak{g})$ as a star-product quantisation of $S(\mathfrak{g})$. Here $U_\hbar(\mathfrak{g})$ denotes the tensor algebra on $\mathfrak{g}$ with relations $vw - wv = \hbar[v, w]$ in terms of the Lie bracket of $\mathfrak{g}$. This is a deformation of $S(\mathfrak{g})$ by the linear map linear map $\varphi : S(\mathfrak{g}) \rightarrow U_\hbar(\mathfrak{g})$ given by a sum over permutations

$$\varphi(v_1 \dots v_n) = \frac{1}{n!} \sum_{\kappa \in S_n} v_{\kappa(1)} \dots v_{\kappa(n)} .$$

As explained in [G], φ is a 1-1 correspondence, and we define a deformed multiplication $\bullet$ on $S(\mathfrak{g})$ by $\underline{v} \bullet \underline{w} = \varphi^{-1}(\varphi(\underline{v}) \cdot \varphi(\underline{w}))$. As examples,

$$\begin{aligned} \varphi(v) \cdot \varphi(w) &= (vw + wv)/2 + \hbar[v, w]/2 \\ &= \varphi(vw) + \hbar \varphi([v, w])/2 , \\ \varphi(v^2) \cdot \varphi(w) &= (v^2w + vwv + wv^2)/3 + \hbar(2v[v, w] + [v, w]v)/3 \\ &= \varphi(v^2w) + \hbar \varphi(v[v, w])/2 + \hbar^2 \varphi([v[v, w]])/6 . \end{aligned}$$

This is related to the CBH formula as follows: since e^v in $S(\mathfrak{g})$ maps under ϕ to e^v in $U_\hbar(\mathfrak{g})$ (similarly for any power series in v) we have

$$e^v \bullet e^w = \phi^{-1}(\phi(e^v)\phi(e^w)) = \phi^{-1}(e^v e^w) = \phi^{-1}(e^{C_\hbar(v, w)}) = e^{C_\hbar(v, w)}$$

where $C_\hbar(v, w)$ is the CBH power series for the product of two exponentials in $U(\mathfrak{g})$ with the insertion of powers of $\hbar$ for each commutator in the Lie algebra.

Lemma 5.2.1. *For $v_1 \dots v_n$ a symmetric product of elements of $\mathfrak{g}$, and $w \in \mathfrak{g}$,*

$$\begin{aligned} w \bullet v_1 \dots v_n &= \varphi(wv_1 \dots v_n) - \hbar n \varphi([v_1, w]v_2 \dots v_n)/2 \\ &\quad + \hbar^2 n(n-1) \varphi([v_1, [v_2, w]]v_3 \dots v_n)/12 + O(\hbar^3) . \end{aligned}$$

Proof: Set

$$\begin{aligned} 0 \leq i \leq n & \quad a_i^0 = v_1 \dots v_i w \dots v_n , \\ 0 \leq i \leq n-1 & \quad a_i^1 = v_1 \dots v_i [v_{i+1}, w] \dots v_n , \\ 0 \leq i \leq n-2 & \quad a_i^2 = v_1 \dots v_i [v_{i+1}, [v_{i+2}, w]] \dots v_n . \end{aligned}$$

Then in $U_\hbar(\mathfrak{g})$ we have

$$a_i^m - a_{i+1}^m = -\hbar a_i^{m+1} ,$$

and this gives, for $i > 0$,

$$(17) \quad a_i^m = a_0^m + \hbar(a_0^{m+1} + \dots + a_{i-1}^{m+1}) .$$

From this we get

$$\begin{aligned} a_0^0 + \dots + a_n^0 &= (n+1)a_0^0 + \hbar(na_0^1 + (n-1)a_1^1 + \dots + a_{n-1}^1) \\ &= (n+1)a_0^0 + \hbar(n+1)(a_0^1 + \dots + a_{n-1}^1)/2 \\ (18) \quad &\quad + \hbar((n-1)a_0^1 + (n-3)a_1^1 + \dots + (1-n)a_{n-1}^1)/2 . \end{aligned}$$

Now we use (17) again to get

$$\begin{aligned}
\sum_{j:0 \leq j \leq n-1} (n-1-2j) a_j^1 &= \hbar \sum_{j:0 \leq j \leq n-1} (n-1-2j) \sum_{i:0 \leq i \leq j-1} a_i^2 \\
&= \hbar \sum_{i:0 \leq i \leq n-2} a_i^2 \sum_{j:i+1 \leq j \leq n-1} (n-1-2j) \\
&= -\hbar \sum_{i:0 \leq i \leq n-2} a_i^2 (i+1)(n-1-i),
\end{aligned}$$

and from this (18) becomes the following, where we use (17) again to get the last equality:

$$\begin{aligned}
a_0^0 + \dots + a_n^0 &= (n+1)a_0^0 + \hbar(n+1)(a_0^1 + \dots + a_{n-1}^1)/2 \\
&\quad - \frac{\hbar^2}{2} \sum_{i:0 \leq i \leq n-2} a_i^2 (i+1)(n-1-i) \\
&= (n+1)a_0^0 + \hbar(n+1)(a_0^1 + \dots + a_{n-1}^1)/2 \\
&\quad - \frac{n(n+1)\hbar^2}{12} (a_0^2 + \dots + a_{n-2}^2) + O(\hbar^3)
\end{aligned} \tag{19}$$

as required. $\heartsuit$

Lemma 5.2.2. *For $v_1 \dots v_n$ and $w_0 \dots w_m$ symmetric products of elements of $\mathfrak{g}$,*

$$\begin{aligned}
w_0 \bullet \varphi(w_1 \dots w_m v_1 \dots v_n) &= \varphi(w_0 w_1 \dots w_m v_1 \dots v_n) \\
&\quad - \hbar n \varphi([v_1, w_0] w_1 \dots w_m v_2 \dots v_n)/2 \\
&\quad + \hbar^2 \varphi(n(n-1)[v_1, [v_2, w_0]] w_1 \dots w_m v_3 \dots v_n \\
&\quad + nm[w_1, [v_1, w_0]] w_2 \dots w_m v_2 \dots v_n)/12 + O(\hbar^3) \\
w_0 \bullet \varphi([v_1, w_1] w_2 \dots w_m v_2 \dots v_n) &= \varphi([v_1, w_0] w_1 \dots w_m v_2 \dots v_n) \\
&\quad - \hbar \varphi([w_0, [w_1, v_1]] w_2 \dots w_m v_2 \dots v_n) \\
&\quad + (n-1)[v_1, w_0][v_2, w_1] w_2 \dots w_m v_3 \dots v_n)/2 + O(\hbar^2).
\end{aligned}$$

Proof: Using 5.2.1 and being careful about counting permutations, we get, for $u \in \mathfrak{g}$, to $O(\hbar^3)$,

$$\begin{aligned}
u \bullet \varphi(w_1 \dots w_m v_1 \dots v_n) &= \varphi(u w_1 \dots w_m v_1 \dots v_n) \\
&\quad - \hbar(n+m) n \varphi([v_1, u] w_1 \dots w_m v_2 \dots v_n)/(2(n+m)) \\
&\quad - \hbar(n+m) m \varphi([w_1, u] w_2 \dots w_m v_1 \dots v_n)/(2(n+m)) \\
&\quad + \hbar^2(n+m)(n+m-1)(1/12) \varphi(\\
&\quad n(n-1)[v_1, [v_2, u]] w_1 \dots w_m v_3 \dots v_n/((n+m)(n+m-1)) \\
&\quad + nm[v_1, [w_1, u]] w_2 \dots w_m v_2 \dots v_n/((n+m)(n+m-1)) \\
&\quad + nm[w_1, [v_1, u]] w_2 \dots w_m v_2 \dots v_n/((n+m)(n+m-1)) \\
&\quad + m(m-1)[w_1, [w_2, u]] w_3 \dots w_m v_1 \dots v_n/((n+m)(n+m-1))) .
\end{aligned}$$

Putting $u = w_0$ and supposing that $w_0 \dots w_m$ is symmetrised, this reduces to the first part of the statement. Next, to $O(\hbar^2)$,

$$\begin{aligned}
u \bullet \varphi([v_1, w_1] w_2 \dots w_m v_2 \dots v_n) &= \varphi(u[v_1, w_1] w_2 \dots w_m v_2 \dots v_n) \\
&\quad - \hbar(n+m-1)(1/2) \varphi(\\
&\quad [[v_1, w_1], u] w_2 \dots w_m v_2 \dots v_n/(n+m-1) \\
&\quad + (m-1)[w_2, u][v_1, w_1] w_3 \dots w_m v_2 \dots v_n/(n+m-1) \\
&\quad + (n-1)[v_2, u][v_1, w_1] w_2 \dots w_m v_3 \dots v_n/(n+m-1)) .
\end{aligned}$$

If $w_0 \dots w_m$ is symmetrised, this reduces to the second part of the statement. $\heartsuit$

Proposition 5.2.3. *For symmetric $w_1 \dots w_m$ and $v_1 \dots v_n$:*

$$\begin{aligned} w_1 \dots w_m \bullet v_1 \dots v_n &= \varphi(w_1 \dots w_m v_1 \dots v_n) \\ &\quad - \hbar m n \varphi([v_1, w_1] w_2 \dots w_m v_2 \dots v_n)/2 \\ &\quad + \hbar^2 n(n-1)m(m-1) \varphi([v_1, w_1][v_2, w_2] w_3 \dots w_m v_3 \dots v_n)/8 \\ &\quad + \hbar^2 n(n-1)m \varphi([v_1, [v_2, w_1]] w_2 \dots w_m v_3 \dots v_n)/12 \\ &\quad + \hbar^2 m(m-1)n \varphi([w_1, [w_2, v_1]] w_3 \dots w_m v_2 \dots v_n)/12 + O(\hbar^3). \end{aligned}$$

Proof: By induction on m . We will suppose that, for fixed n and symmetric $w_1 \dots w_m$,

$$\begin{aligned} w_1 \dots w_m \bullet v_1 \dots v_n &= \varphi(w_1 \dots w_m v_1 \dots v_n) - \hbar \alpha_m \varphi([v_1, w_1] w_2 \dots w_m v_2 \dots v_n) \\ &\quad + \hbar^2 (\beta_m \varphi([v_1, w_1][v_2, w_2] w_3 \dots w_m v_3 \dots v_n) \\ &\quad + \gamma_m \varphi([v_1, [v_2, w_1]] w_2 \dots w_m v_3 \dots v_n) \\ &\quad + \delta_m \varphi([w_1, [w_2, v_1]] w_3 \dots w_m v_2 \dots v_n)) + O(\hbar^3). \end{aligned}$$

Using associativity of the $\bullet$ product,

$$\begin{aligned} w_0 \dots w_m \bullet v_1 \dots v_n &= \varphi(w_0 w_1 \dots w_m v_1 \dots v_n) \\ &\quad - \hbar n \varphi([v_1, w_0] w_1 \dots w_m v_2 \dots v_n)/2 \\ &\quad + \hbar^2 \varphi(n(n-1)[v_1, [v_2, w_0]] w_1 \dots w_m v_3 \dots v_n \\ &\quad - nm[w_0, [w_1, v_1]] w_2 \dots w_m v_2 \dots v_n)/12 \\ &\quad - \hbar \alpha_m \varphi([v_1, w_0] w_1 \dots w_m v_2 \dots v_n) \\ &\quad + \hbar^2 \alpha_m \varphi([w_0, [w_1, v_1]] w_2 \dots w_m v_2 \dots v_n \\ &\quad + (n-1)[v_1, w_0][v_2, w_1] w_2 \dots w_m v_3 \dots v_n)/2 \\ &\quad + \hbar^2 (\beta_m \varphi([v_1, w_0][v_2, w_1] w_2 \dots w_m v_3 \dots v_n) \\ &\quad + \gamma_m \varphi([v_1, [v_2, w_0]] w_1 \dots w_m v_3 \dots v_n) \\ &\quad + \delta_m \varphi([w_0, [w_1, v_1]] w_2 \dots w_m v_2 \dots v_n)) + O(\hbar^3). \end{aligned}$$

This gives the recursive equations and initial conditions

$$\begin{aligned} \alpha_{m+1} &= \alpha_m + n/2, \quad \alpha_1 = n/2, \\ \beta_{m+1} &= \beta_m + \alpha_m(n-1)/2, \quad \beta_1 = 0, \\ \gamma_{m+1} &= \gamma_m + n(n-1)/12, \quad \gamma_1 = n(n-1)/12, \\ \delta_{m+1} &= \delta_m - nm/12 + \alpha_m/2, \quad \delta_1 = 0. \end{aligned}$$

From this we get $\alpha_m = nm/2$, $\gamma_m = n(n-1)m/12$, $\delta_m = m(m-1)n/12$ and $\beta_m = n(n-1)m(m-1)/8$. $\heartsuit$

5.3. Cochain for the deformed product of $S(\mathfrak{g})$. Choose a dual basis $e_i \in \mathfrak{g}$ and $e^i \in \mathfrak{g}^*$. Let $\mathfrak{g}$ act on $S(\mathfrak{g})$ by the adjoint, and $\mathfrak{g}^*$ act by evaluation. Set $Q_1 = e_i \otimes e^i$ and $Q_2 = e^i \otimes e_i$, and let μ stand for multiplication. Then for symmetric $\underline{w} = w_1 \dots w_m$ and $\underline{v} = v_1 \dots v_n$:

$$\begin{aligned} \mu(e_i \otimes e^i)(\underline{w} \otimes \underline{v}) &= n[v_1, w_1 \dots w_m] v_2 \dots v_n \\ &= nm[v_1, w_1] w_2 \dots w_m v_2 \dots v_n \\ \mu(e^i \otimes e_i)(\underline{w} \otimes \underline{v}) &= m w_2 \dots w_m [w_1, v_1 \dots v_n] \\ &= -nm[v_1, w_1] w_2 \dots w_m v_2 \dots v_n. \end{aligned}$$

We have quadratic terms of the form Q_1^2 , Q_2^2 and $:Q_1 Q_2:$ (with $::$ denoting a normal ordering with elements of $\mathfrak{g}^*$ being put on the right), which are respectively

$$\begin{aligned} \mu(e_i e_j \otimes e^i e^j)(\underline{w} \otimes \underline{v}) &= n(n-1)[v_1, [v_2, w_1 \dots w_m]] v_3 \dots v_n \\ &= n(n-1)m[v_2, [v_1, w_1] w_2 \dots w_m] v_3 \dots v_n \\ &= n(n-1)m[v_2, [v_1, w_1]] w_2 \dots w_m v_3 \dots v_n \\ &\quad + n(n-1)m(m-1)[v_1, w_1][v_2, w_2] w_3 \dots w_m v_3 \dots v_n, \\ \mu(e^i e^j \otimes e_i e_j)(\underline{w} \otimes \underline{v}) &= m(m-1) w_3 \dots w_m [w_1, [w_2, v_1 \dots v_n]] \end{aligned}$$

$$\begin{aligned}
&= nm(m-1) w_3 \dots w_m [w_1, [w_2, v_1] v_2 \dots v_n] \\
&= nm(m-1) w_3 \dots w_m [w_1, [w_2, v_1]] v_2 \dots v_n \\
&\quad + n(n-1)m(m-1) w_3 \dots w_m [w_2, v_2] [w_1, v_1] v_3 \dots v_n, \\
\mu(e_i e^j \otimes e_j e^i)(\underline{w} \otimes \underline{v}) &= n[v_1, e^j(\underline{w})] \cdot e_j(v_2 \dots v_n) \\
&= nm[v_1, w_2 \dots w_m] \cdot [w_1, v_2 \dots v_n] \\
&= -n(n-1)m(m-1) [v_1, w_1] [v_2, w_2] w_3 \dots w_m v_3 \dots v_n.
\end{aligned}$$

If we set

$$(20) \quad F^{-1} = 1 \otimes 1 + \hbar(\alpha Q_1 + \beta Q_2) + \hbar^2(2Q_1^2 + 2Q_2^2 + :Q_1 Q_2:)/24 + O(\hbar^3),$$

where $\alpha - \beta = -1/2$, then we recover the CBH product to $O(\hbar^3)$. Note that we could add any multiple of $Q_1 Q_2 + Q_2^2$ or $Q_2 Q_1 + Q_1^2$ to $G^{(2)}$ and still get the same product to $O(\hbar^3)$. Also we have the equation

$$\mu(:Q_1 Q_2:^R + :Q_1 Q_2: + Q_1^2 + Q_2^2)(\underline{w} \otimes \underline{v}) = nm w_2 \dots w_m v_2 \dots v_n \cdot e^i([v_1, [w_1, e_i]]) ,$$

where $:Q_1 Q_2:^R = e^j e_i \otimes e^i e_j$ is the reversed normal order. Now $e^i([v, [w, e_i]])$ (summed over i) is the trace of $\text{ad}_v \text{ad}_w$, that is $-\langle v, w \rangle$, where $\langle, \rangle$ is the killing form. If we set $\langle e_i, e_j \rangle = \kappa_{ij}$, then

$$\mu(:Q_1 Q_2:^R + :Q_1 Q_2: + Q_1^2 + Q_2^2 + \kappa_{ij} e^i \otimes e^j)(\underline{w} \otimes \underline{v}) = 0.$$

5.4. Improved cochain for the deformed coproduct on $S(\mathfrak{g}^*)$. Here we consider a more general covariant ansatz but show that a further requirement relating to the coproduct of $S(\mathfrak{g})$ again leads to a unique answer. Thus, we can write a more general expression for $G^{(2)}$ as

$$(21) \quad G^{(2)} = (2Q_1^2 + 2Q_2^2 + :Q_1 Q_2:)/24 + \gamma(Q_1 + Q_2)Q_1 + \delta(Q_1 + Q_2)Q_2 \\ + \zeta(:Q_1 Q_2:^R + :Q_1 Q_2: + Q_1^2 + Q_2^2 + \kappa_{ij} e^i \otimes e^j),$$

where γ, δ and ζ are constants, and we shall use this instead of the $\hbar^2$ term in (20). By the above discussion, this still gives the correct deformed product on $S(\mathfrak{g})$. We use $G^{(1)} = \alpha Q_1 + \beta Q_2$. Then from (3) the deformed coproduct is given by

$$(22) \quad \Delta_F(x) = \Delta(x) + \hbar[\Delta(x), G^{(1)}] + \hbar^2([\Delta(x), G^{(2)}] - G^{(1)}[\Delta(x), G^{(1)}]) + O(\hbar^3).$$

For $x \in \mathfrak{g}^*$ we have $\Delta(x) = x \otimes 1 + 1 \otimes x$, and using the fact that elements of $\mathfrak{g}^*$ commute, we find

$$[\Delta(x), G^{(1)}] = \alpha[x, e_i] \otimes e^i + \beta e^i \otimes [x, e_i].$$

The coefficient of $\hbar^2$ in (22) is

$$\begin{aligned}
&\left(\gamma[x, e_i]e_j + (\gamma - \alpha^2)e_j[x, e_i] + \left(\frac{1}{12} + \zeta\right)[[x, e_i], e_j] \right) \otimes e^i e^j \\
&+ e^i e^j \otimes \left(\delta[x, e_i]e_j + (\delta - \beta^2)e_j[x, e_i] + \left(\frac{1}{12} + \zeta\right)[[x, e_i], e_j] \right) \\
&+ e^i[x, e_j] \otimes \left(\left(\gamma + \frac{1}{24} - \alpha\beta + \zeta\right)e_i e^j + (\delta + \zeta)e^j e_i \right) \\
&+ \left((\gamma + \zeta)e^i e_j + \left(\delta + \frac{1}{24} - \alpha\beta + \zeta\right)e_j e^i \right) \otimes [x, e_i] e^j.
\end{aligned}$$

To ensure that this is in $S(\mathfrak{g}^*) \otimes S(\mathfrak{g}^*)$ we require that $\gamma = \alpha^2/2$, $\delta = \beta^2/2$ and $(\alpha - \beta)^2 = -4\zeta - 1/12$. We already have $\alpha - \beta = -1/2$, so we get $\zeta = -1/12$. Now the coefficient of $\hbar^2$ in (22) is:

$$\begin{aligned}
\Delta_F(x) &= x \otimes 1 + 1 \otimes x + \hbar(\alpha[x, e_i] \otimes e^i + \beta e^i \otimes [x, e_i]) \\
&\quad + \hbar^2 \alpha^2 [[x, e_i], e_j] \otimes e^i e^j / 2 + e^i e^j \otimes \beta^2 [[x, e_i], e_j] / 2 \\
&\quad + (\beta^2/2 - 1/12) e^i[x, e_j] \otimes [e^j, e_i] + (\alpha^2/2 - 1/12) [e^i, e_j] \otimes [x, e_i] e^j.
\end{aligned}$$

Also putting these values into the general form of F^{-1} we obtain:

Theorem 5.4.1. *The cochain*

$$F^{-1} = 1 \otimes 1 + \hbar(\alpha Q_1 + \beta Q_2) + \hbar^2 \left(\frac{\alpha^2}{2} (Q_1 + Q_2) Q_1 + \frac{\beta^2}{2} (Q_1 + Q_2) Q_2 - \frac{1}{24} (: Q_1 Q_2 : + 2 : Q_1 Q_2 :^R + 2\kappa_{ij} e^i \otimes e^j) \right) + O(\hbar^3)$$

for $\alpha - \beta = -\frac{1}{2}$ reproduces the product of $U_\hbar(\mathfrak{g})$ to the relevant order and has the property that $\Delta_F(S(\mathfrak{g}^*)) \subset S(\mathfrak{g}^*) \otimes S(\mathfrak{g}^*)$ to the relevant order.

It appears that the up to the choice of how $-\frac{1}{2}$ is split between α and β , the cochain F is determined at higher orders by the properties in the theorem to hold for any Lie algebra and the requirement of having a $\mathfrak{g}$ -invariant form. Although we will not give a formal proof, let us explain the underlying reason here. First we note that locally near the identity we may identify the Lie algebra with the connected and simply connected Lie group G associated to it, i.e. $S(\mathfrak{g}^*)_F \subseteq \mathbb{C}_{loc}[G]$, where the latter denotes functions defined near the identity. This is by

$$\Theta : S(\mathfrak{g}^*) \supset \mathfrak{g}^* \ni x \mapsto f_x(e^{\hbar v}) = \hbar^{-1} \langle x, v \rangle$$

(i.e. the generators appear as logarithmic coordinates on the Lie group). Let us now show that

$$((\Theta \otimes \Theta) \Delta_F x)(e^v, e^w) = (\Delta_{\mathbb{C}_{loc}[G]} \Theta(x))(e^v, e^w),$$

i.e. the identification is indeed as Hopf algebras to the relevant order. The right hand side here is $\Theta(x)(e^v e^w) = \Theta(x)(e^{C(v,w)}) = \hbar^{-1} \langle x, C(v,w) \rangle$ where $C(v,w)$ is the Campbell-Baker-Hausdorff series. Meanwhile, evaluating the coproduct Δ_F we have

$$\begin{aligned} ((\Theta \otimes \Theta) \Delta_F x)(e^v, e^w) &= \hbar^{-1} (\langle x, v \rangle + \langle x, w \rangle) + \hbar^{-1} \alpha \langle [x, e_i], v \rangle \langle e^i, w \rangle + \hbar^{-1} \beta \langle e^i, v \rangle \langle [x, e_i], w \rangle \\ &\quad + \hbar^{-1} \frac{\alpha^2}{2} \langle v, [[x, e_i], e_j] \rangle \langle e^i, w \rangle \langle e^j, w \rangle + \hbar^{-1} \frac{\beta^2}{2} \langle w, e^i \rangle \langle w, e^j \rangle \langle v, [[x, e_i], e_j] \rangle \\ &\quad + \hbar^{-1} \left(\frac{\alpha^2}{2} - \frac{1}{12} \right) \langle v, e^i \rangle \langle v, [x, e_j] \rangle \langle w, [e^j, e_i] \rangle + \hbar^{-1} \left(\frac{\beta^2}{2} - \frac{1}{12} \right) \langle v, e^i \rangle \langle v, e^j \rangle \langle w, [[x, e_i], e_j] \rangle + \dots \\ &= \hbar^{-1} \langle x, v + w + [w, v] (\alpha - \beta) + \frac{\alpha^2}{2} [w, [w, v]] + \frac{\beta^2}{2} [v, [v, w]] + \left(\frac{\alpha^2}{2} - \frac{1}{12} \right) [[w, v], w] \\ &\quad + \left(\frac{\beta^2}{2} - \frac{1}{12} \right) [[v, w], v] + \dots \\ &= \hbar^{-1} \langle x, v + w + \frac{1}{2} [v, w] + \frac{1}{12} ([v, [v, w]] + [[v, w], w]) + \dots \rangle \end{aligned}$$

for all $x \in \mathfrak{g}^*$. We used $\langle [x, v], w \rangle = \langle x \triangleleft v, w \rangle = \langle x, v \triangleright w \rangle = \langle x, [v, w] \rangle$ and similarly for repeated commutators. Note that the increasing powers of $\hbar^{-1}$ with each evaluation exactly match the increasing powers of $\hbar$ in the powerseries coming from F . Hence the twisted coproduct reproduces the Campbell-Baker-Hausdorff series to low degree in its expansion. It is clear that this requirement and that we continue to reproduce the product of $U_\hbar(\mathfrak{g})$ can be used to determine a universal formula for F, F^{-1} , though again it would be beyond our scope to provide this here.

5.5. The Duflo map. The Duflo map provides an independent check of our formula in Theorem 5.4.1 and gives some idea of the structure of F^{-1} at all orders. We recall [Du] that there is an invertible operator D on $S(\mathfrak{g})$ defined by

$$D = e^{\sum_{k=1}^{\infty} \alpha_{2k} \partial_{\text{Tr}_{2k}}}; \quad \sum_{k=1}^{\infty} \alpha_{2k} t^{2k} \equiv \frac{1}{2} \ln \left(\frac{\sinh(t/2)}{t/2} \right)$$

where $\partial_{\text{Tr}_{2k}}$ is the differential operator on $S(\mathfrak{g})$ given by the action of the element $\text{Tr}_{2k} = \text{Tr}_{\mathfrak{g}}((\text{ad.})^{2k}) \in S(\mathfrak{g}^*)$. The lowest order part is

$$\alpha_2 = \frac{1}{48}, \quad \alpha_4 = -\frac{1}{5760}, \quad \partial_{\text{Tr}_2} = -\kappa_{ij} e^i e^j$$

in our conventions above. Duflo's theorem is that when restricted to the ad-invariant subalgebra $S(\mathfrak{g})^{\mathfrak{g}}$ the map $\varphi \circ D$ is an isomorphism of this with the centre $Z(U(\mathfrak{g}))$.

Now let $F_{\text{red}}^{-1} \in U(\mathfrak{g}^*)^{\otimes 2}$ be the effective F^{-1} when acting on invariants $S(\mathfrak{g})^{\mathfrak{g}} \otimes S(\mathfrak{g})^{\mathfrak{g}}$. This is given by normal ordering F^{-1} so that all terms have all elements of $\mathfrak{g}$ to the right of all elements of $\mathfrak{g}^*$ (what we called $: :^R$ above). Then project $\mathfrak{g}$ to zero in the result because by definition it act by zero on invariant elements. The result is some power-series in $U(\mathfrak{g}^*)^{\otimes 2} = S(\mathfrak{g}^*)^{\otimes 2}$ since $\mathfrak{g}^*$ is being regarded as an Abelian Lie algebra.

Proposition 5.5.1. *At least to $O(\hbar^3)$, F_{red}^{-1} for the cochain in Theorem 5.4.1 is a coboundary in the sense of [M2, Chapter 2.3] of the Duflo element, i.e.,*

$$F_{\text{red}}^{-1} = (\Delta\gamma)(\gamma^{-1} \otimes \gamma^{-1}), \quad \gamma = e^{\sum_{k=1}^{\infty} \alpha_{2k} \hbar^{2k} \text{Tr}_{2k}}$$

Here γ viewed as an operator acting on $S(\mathfrak{g})$ is just D in Duflo's theorem after explicitly introducing the deformation scaling parameter. The coproduct Δ is that of $U(\mathfrak{g}^*)$. We expect this result to hold to all orders because F_{red}^{-1} a coboundary of some cochain γ implies that

$$f \bullet g = \mu(F_{\text{red}}^{-1} \cdot (f \otimes g)) = \gamma \cdot \mu(\gamma^{-1} \cdot f \otimes \gamma^{-1} \cdot g) = D(\mu(D^{-1} f \otimes D^{-1} g))$$

for all $f, g \in S(\mathfrak{g})^{\mathfrak{g}}$, where D denotes γ acting on $S(\mathfrak{g})$. We used in the first equality that $U(\mathfrak{g}^*)$ acts covariant on $S(\mathfrak{g})$ with its initial product μ and hence we can move the action of $\Delta\gamma$ to the left as the action of γ . This means that the modified product restricted to invariant elements is an isomorphism of algebras (this is the meaning of F_{red}^{-1} being a coboundary as explained in [M2]). In the light of Duflo's theorem we expect F_{red}^{-1} therefore to be a coboundary of an invertible element γ whose action is the same as the operator D in Duflo's theorem. This leads to the statement of the proposition.

We now verify the proposition to the order $O(\hbar^3)$ available to us. In the expression in Theorem 5.4.1 all the terms have some e_i already to the right and therefore fail to contribute, except $: Q_1 Q_2 :$ and the $\kappa_{ij} e^i \otimes e^j$ terms. We write the former using

$$e_j e^i = e^i e_j + (e^i \triangleleft e_j) = e^i e_j + f_{jki} e^k$$

where $[e_i, e_j] = f_{ijk} e_k$ defines the structure constants. Then

$$: Q_1 Q_2 := e_i e^j \otimes e_j e^i \sim f_{jki} f_{imj} e^k \otimes e^m \sim f_{kji} f_{mij} e^k \otimes e^m \sim -\kappa_{ij} e^i \otimes e^j$$

discarding terms acting trivially on invariant elements. As a result we have

$$F_{\text{red}}^{-1} = 1 \otimes 1 - \frac{\hbar^2}{24} \kappa_{ij} e^i \otimes e^j + O(\hbar^3)$$

(in fact the next term should be $O(\hbar^4)$). On the other hand from the above

$$\gamma = e^{-\frac{\hbar^2}{48} c + O(\hbar^4)}; \quad c = \kappa_{ij} e^i e^j, \quad (\Delta\gamma)(\gamma^{-1} \otimes \gamma^{-1}) = e^{-\frac{\hbar^2}{48} (\Delta c - c \otimes 1 - 1 \otimes c) + O(\hbar^4)} = F_{\text{red}}^{-1}$$

to lowest order. The same proposition would provide a check to all orders of any cochain found. At the moment we have provided a check of our order $O(\hbar^3)$ result.

5.6. Example: noncommutative Minkowski space as cochain twist. Here we verify (20) for the algebra $[t, x_i] = \hbar x_i$ which has been proposed as noncommutative spacetime (the so-called bicrossproduct model). For convenience we take only one $x = x_i$ rather than $i = 1, 2, 3$ for spacetime, however the structure is exactly similar. In this case we exhibit a candidate for F^{-1} to the next order, i.e. up to $O(\hbar^4)$.

In this model there is a representation of the algebra in terms of 2×2 matrices, as

$$t \mapsto \begin{pmatrix} \hbar/2 & 0 \\ 0 & -\hbar/2 \end{pmatrix}, \quad x \mapsto \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

These matrices can be exponentiated to give

$$\exp(pt + qx) = \begin{pmatrix} e^{\frac{\hbar p}{2}} & \frac{(-1+e^{\hbar p})q}{e^{\frac{\hbar p}{2}} - \hbar p} \\ 0 & e^{-\frac{\hbar p}{2}} \end{pmatrix}.$$

A little matrix multiplication shows that

$$\exp(pt + qx) \cdot \exp(rt + sx) = \exp\left((p+r)t + \frac{(p+r) \left((-1+e^{\hbar p})qr + e^{\hbar p}(-1+e^{\hbar r})ps\right)}{(-1+e^{\hbar(p+r)})pr}\right)x,$$

so in this case we have a closed form for the CBH formula.

Further calculation with this algebra gives

$$\begin{aligned} \varphi(x^n t^m) &= x^n \left(t^m + \frac{nm}{2} \hbar t^{m-1} + \frac{n(3n+1)m(m-1)}{24} \hbar^2 t^{m-2} \right. \\ &\quad \left. + \frac{n^2(n+1)m(m-1)(m-2)}{48} \hbar^3 t^{m-3} + O(\hbar^4) \right). \end{aligned}$$

From this we can calculate

$$\begin{aligned} \varphi^{-1}(x^n t^m) &= x^n \left(t^m - \frac{nm}{2} \hbar t^{m-1} + \frac{n(3n-1)m(m-1)}{24} \hbar^2 t^{m-2} \right. \\ &\quad \left. + \frac{n^2(1-n)m(m-1)(m-2)}{48} \hbar^3 t^{m-3} + O(\hbar^4) \right). \end{aligned}$$

If we combine this with the following formula for multiplication in $U_\hbar$,

$$t^s x^r = \sum_{p=0}^s C_p^s (r \hbar)^p x^r t^{s-p},$$

(where C_p^s is a binomial coefficient) we get the formula

$$\begin{aligned} (x^n t^m) \bullet (x^r t^s) &= x^{n+r} t^{m+s} + \frac{\hbar}{2} (mr - ns) x^{n+r} t^{m+s-1} \\ &\quad + \frac{\hbar^2}{24} (mr - m^2 r - 3mr^2 + 3m^2 r^2 + ns - 2mns - 3n^2 s - 2mrs - 6mnrs \\ &\quad - ns^2 + 3n^2 s^2) x^{n+r} t^{m+s-2} + \frac{\hbar^3}{48} (-2mr^2 + 3m^2 r^2 - m^3 r^2 + 2mr^3 \\ &\quad - 3m^2 r^3 + m^3 r^3 + 2n^2 s - 2mn^2 s - 2n^3 s - m^2 nrs - 3mn^2 rs + 2mr^2 s \\ &\quad - 2m^2 r^2 s + 3mnr^2 s - 3m^2 nr^2 s - 3n^2 s^2 + 2mn^2 s^2 + 3n^3 s^2 + mnrs^2 \\ &\quad + 3mn^2 rs^2 + n^2 s^3 - n^3 s^3) x^{n+r} t^{m+s-3} + O(\hbar^4). \end{aligned} \tag{23}$$

On $S(\mathfrak{g})$, ad_t is identified with $x \frac{d}{dx}$, and ad_x is identified with $-x \frac{d}{dt}$. We take the dual basis $\hat{t}, \hat{x} \in \mathfrak{g}^*$. Then on $S(\mathfrak{g})$, $\hat{t}$ is identified with $\frac{d}{dt}$, and $\hat{x}$ is identified with $\frac{d}{dx}$. Using this, it can be

explicitly checked that (20) gives the deformed multiplication for this algebra up to $O(\hbar^3)$. The third order part of (23) can be given by

$$G^{(3)} = (e_i e^j e^k \otimes e_k e_j e^i - e_k e^j e_i \otimes e^i e_j e^k - 2 e^i e_j e^k \otimes e_i e^j e_k) / 96 ,$$

where we sum over i, j, k . Note, however, that this expression is not unique in the same manner that (20) at order $\hbar^2$ is not unique as we have seen. With more work one may exploit the non-uniqueness and expect to achieve the features in Section 5.4 with respect to the coproduct Δ_F as well. Note that we do not necessarily expect a unique F, F^{-1} for any given Lie algebra (the uniqueness proposed in Section 5.4 was for a universal F, F^{-1} applicable to all Lie algebras).

6. MACKEY QUANTISATIONS $C^\infty(N) \rtimes U_\hbar(\mathfrak{g})$ OF HOMOGENEOUS SPACES AS COCHAIN TWISTS

Suppose that a Lie group G with Lie algebra $\mathfrak{g}$ acts on a manifold N . In this case there is a standard 'quantisation' for the system due to Mackey and much used in physics, in which the initial algebra is $C^\infty(N) \otimes S(\mathfrak{g}) \subset C^\infty(N \times \mathfrak{g}^*)$ (i.e. functions polynomial in the $\mathfrak{g}^*$ direction). This is deformed or quantised to the cross product $C^\infty(N) \rtimes U_\hbar(\mathfrak{g})$. Here, for $v \in \mathfrak{g}$ and $f \in C^\infty(N)$, $(v \triangleright_\hbar f)(x) = \hbar f'(x; v(x))$, and $U_\hbar(\mathfrak{g})$ has the relation $vw - wv = \hbar[v, w]$ in terms of the Lie bracket $[\cdot, \cdot]$ on $\mathfrak{g}$. This algebra acts on the L^2 sections of a bundle whose fiber over $x \in N$ is a representation of the stabiliser of x in G . In this section we show that the results of Section 5 may be extended to this case also. The theory here reduces to that of Section 5 when N is a point.

Note that $M = N \times \mathfrak{g}^*$ is indeed a Poisson manifold, because the quantisation above can be viewed as a flat deformation. Its Poisson bracket has a semidirect product form

$$\{f, g\} = 0, \quad \{v, f\} = v \triangleright f, \quad \{v, w\} = [v, w]$$

for $f, g \in C^\infty(N)$ and $v, w \in \mathfrak{g}^*$. Our goal is to lift this Poisson bivector to an element of a suitable $\mathcal{L} \otimes \mathcal{L}$ and hence to a cochain F at least to order $O(\hbar^2)$, i.e. to express the Mackey quantisation as a cochain twist.

6.1. To first order. We will be extending the results from the previous 'CBH' case in Section 5; we denote the cochain components there by $G_{CBH}^{(i)}$.

Definition 6.1.1. Take a dual basis (e_i, e^i) with $e_i \in \mathfrak{g}$ and $e^i \in \mathfrak{g}^*$. Then define some vector fields on $M = N \times \mathfrak{g}^*$ by the following formulae, where $v \in \mathfrak{g} \subset S(\mathfrak{g})$ and $g \in C^\infty(N)$.

$$\begin{aligned} \check{e}_i(v) &= [e_i, v] & \check{e}_i(g) &= 0, \\ \check{e}^i(v) &= e^i(v) & \check{e}^i(g) &= 0, \\ \check{c}_i(v) &= 0 & \check{c}_i(g) &= e_i \triangleright g. \end{aligned}$$

Proposition 6.1.2. The Poisson structure described is given by

$$G^{(1)} = -(\check{c}_i + \check{e}_i/2) \otimes \check{e}^i.$$

Proof: Recall that for $G^{(1)} = \sum X \otimes Y$ we have $\{a, b\} = \sum X(a)Y(b) - Y(a)X(b)$. Hence for $v, w \in \mathfrak{g}$ and $g, k \in C^\infty(N)$:

$$\begin{aligned} \check{e}_i(v) \check{e}^i(w) - \check{e}^i(v) \check{e}_i(w) &= [e_i, v] e^i(w) - e^i(v) [e_i, w] = [w, v] - [v, w] = 2[w, v], \\ \check{e}_i(v) \check{e}^i(g) - \check{e}^i(v) \check{e}_i(g) &= 0, \\ \check{e}_i(g) \check{e}^i(k) - \check{e}^i(g) \check{e}_i(k) &= 0, \\ \check{c}_i(v) \check{e}^i(w) - \check{e}^i(v) \check{c}_i(w) &= 0, \\ \check{c}_i(v) \check{e}^i(g) - \check{e}^i(v) \check{c}_i(g) &= -e^i(v) (e_i \triangleright g) = -v \triangleright g, \\ \check{c}_i(g) \check{e}^i(k) - \check{e}^i(g) \check{c}_i(k) &= 0 \end{aligned}$$

as desired. $\heartsuit$

Proposition 6.1.3. *The Lie brackets between the $\check{e}_i$, $\check{c}_i$ and $\check{e}^i$ are given as follows. The $\check{c}_i$ commute with both the $\check{e}_j$ and the $\check{e}^j$. The $\check{e}_i$ have the usual Lie bracket for $\mathfrak{g}$ among themselves. The $\check{e}^i$ commute among themselves. The $\check{c}_i$ have the usual Lie bracket for $\mathfrak{g}$ among themselves. The bracket of the e_i with the e^j is given by the coadjoint action. Thus the given fields form a Lie algebra $\mathcal{L} = \mathfrak{g} \ltimes \mathfrak{g}^* \oplus \mathfrak{g}$.*

Proof: Check against v and g . The most difficult ones are:

$$\begin{aligned} [\check{c}_i, \check{c}_j](g) &= (e_i \triangleright (e_j \triangleright g) - e_j \triangleright (e_i \triangleright g)) = ([e_i, e_j] \triangleright g) , \\ [\check{e}_i, \check{e}^j](v) &= -\check{e}^j([e_i, v]) = -e^j([e_i, v]) . \end{aligned}$$

The last equation shows the coadjoint action, $\text{coad}_w(\psi) = -\psi \circ \text{ad}_w$. $\heartsuit$

We shall use this Lie algebra to induce the Mackey quantisation. We have already seen above that this is sufficient at order $\hbar$.

Proposition 6.1.4. *The preconnection is for $v, w \in \mathfrak{g}$ and $f, g \in C^\infty(N)$:*

$$\begin{aligned} \hat{\nabla}_v dw &= d[v, w]/2 , \\ \hat{\nabla}_v dg &= d(v \triangleright g) , \\ \hat{\nabla}_f dw &= 0 , \\ \hat{\nabla}_f dg &= 0 . \end{aligned}$$

Proof: We use the formula involving Lie derivatives;

$$\hat{\nabla}_a \xi = \check{e}^i(a) \mathcal{L}_{\check{c}_i + \check{e}_i/2} \xi - (\check{c}_i + \check{e}_i/2)(a) \mathcal{L}_{\check{e}^i} \xi .$$

This gives

$$\begin{aligned} \hat{\nabla}_v dw &= e^i(v) d(\check{c}_i + \check{e}_i/2)(w) - [e_i, v] d(\check{e}^i)(w)/2 \\ &= e^i(v) d[e_i, w]/2 , \\ \hat{\nabla}_v dg &= e^i(v) d(\check{c}_i + \check{e}_i/2)(g) - [e_i, v] d(\check{e}^i)(g)/2 \\ &= e^i(v) d(e_i \triangleright g) , \\ \hat{\nabla}_f dw &= -(e_i \triangleright f) d(\check{e}_i)(w) = 0 , \\ \hat{\nabla}_f dg &= -(e_i \triangleright f) d(\check{e}_i)(g) = 0 \end{aligned}$$

as required. $\heartsuit$

Proposition 6.1.5. *The curvature is given by, for $v, w, z \in \mathfrak{g}$ and $f, g, h \in C^\infty(N)$:*

$$\begin{aligned} R(v, w)(dz) &= -d([v, w], z)/4 , \\ R(v, w)(dg) &= 0 , \\ R(v, g)(dz) &= R(v, g)(dh) = 0 , \\ R(f, g)(dz) &= R(f, g)(dh) = 0 . \end{aligned}$$

Proof: First we need to state the Lie brackets of the vector fields, using $[\hat{a}, \hat{b}] = \widehat{\{a, b\}}$:

$$\begin{aligned} [\hat{v}, \hat{w}] &= \widehat{\{v, w\}} = \widehat{[v, w]} , \\ [\hat{v}, \hat{g}] &= \widehat{\{v, g\}} = \widehat{(v \triangleright g)} , \\ [\hat{f}, \hat{g}] &= \widehat{\{f, g\}} = 0 . \end{aligned}$$

Now we find the curvatures. For $R(v, w)(dz)$ the computation is as in Proposition 5.1.1 with the same result. For the new case we have

$$R(v, w)(dg) = d(v \triangleright (w \triangleright g) - w \triangleright (v \triangleright g) - [v, w] \triangleright g) = 0$$

since $\triangleright$ is a representation of the Lie algebra (if it were a cocycle representation we would have curvature). In the expressions for $R(v, g)$ and $R(f, g)$ every term is individually zero. $\heartsuit$

6.2. To second order. The multiplication on $C^\infty(N) \rtimes U_\hbar(\mathfrak{g})$ is given by

$$(f \otimes \underline{v})(g \otimes \underline{w}) = f \cdot (\underline{v}_{(1)} \triangleright_\hbar g) \otimes \underline{v}_{(2)} \bullet \underline{w},$$

where the second product is in $U_\hbar(g)$ and the coproduct is the usual $\Delta v = v \otimes 1 + 1 \otimes v$ for $v \in \mathfrak{g}$. In terms of deformations of $C^\infty(N) \otimes S(\mathfrak{g})$ we can decompose the order $\hbar^2$ part of the product as

$$(24) \quad \begin{aligned} & f \cdot g \otimes (\hbar^2 \text{ part of } \underline{v} \bullet \underline{w}) + f \cdot (\hbar \text{ part of } \underline{v}_{(1)} \triangleright_\hbar g) \otimes (\hbar \text{ part of } \underline{v}_{(2)} \bullet \underline{w}) \\ & + f \cdot (\hbar^2 \text{ part of } \underline{v}_{(1)} \triangleright_\hbar g) \otimes (\hbar^0 \text{ part of } \underline{v}_{(2)} \bullet \underline{w}). \end{aligned}$$

The first term of (24) is given by $\hbar^2 (1 \otimes G_{CBH1}^{(2)}) \otimes (1 \otimes G_{CBH2}^{(2)})$, where the final suffices 1, 2 denote the two pieces of $G_{CBH}^{(2)}$ (summation understood).

The second term of (24) is, where $\hat{}$ denotes omission,

$$\hbar f(x) \cdot \sum_i (v_i \triangleright g)(x) \otimes (\hbar \text{ part of } (v_1 \dots \hat{v}_i \dots v_n) \bullet (w_1 \dots w_m)).$$

We can separate this into two stages, first the moving the v_i stage, and then the $\bullet$ multiplication. The first is given by $\hbar (1 \otimes \check{e}^i) \otimes (\check{e}_i \otimes 1)$, and the second by $\hbar (1 \otimes G_{CBH1}^{(2)}) \otimes (1 \otimes G_{CBH2}^{(2)})$ as above. The third term of (24) is,

$$\hbar^2 f \cdot \sum_{i < j} (v_i v_j \triangleright g) \otimes v_1 \dots \hat{v}_i \dots \hat{v}_j \dots v_n w_1 \dots w_m.$$

This is given by $\frac{1}{2} \hbar^2 (1 \otimes \check{e}^i \check{e}^j) \otimes (1 \otimes \check{e}_i \check{e}_j)$, giving in total

$$G^{(2)} = (1 \otimes G_{CBH1}^{(2)}) \otimes (1 \otimes G_{CBH2}^{(2)}) + \sum_i (1 \otimes G_{CBH1}^{(1)} \check{e}^i) \otimes (\check{e}_i \otimes G_{CBH2}^{(1)}) + \frac{1}{2} \sum_{ij} (1 \otimes \check{e}^i \check{e}^j) \otimes (1 \otimes \check{e}_i \check{e}_j)$$

A special case is of course $N = G$ and action by left translation. Then the Mackey quantisation $C^\infty(G) \rtimes U_\hbar(\mathfrak{g})$ is a quantisation of $T^*G = G \times \mathfrak{g}^*$ and the Poisson-bracket above becomes the standard symplectic structure on T^*G . In that case we have an actual connection ∇ in Section 6.1.

6.3. Special case of T^*G . In general we have a Poisson map $T^*N \rightarrow N \times \mathfrak{g}^*$ defined using the moment map by $(n, p) \mapsto (n, \langle x_-(n), p \rangle)$ where x_ξ is the vector field for the action of $\xi \in \mathfrak{g}$ on N . This means a map

$$C^\infty(N \times \mathfrak{g}^*) \rightarrow C^\infty(T^*N)$$

which will be surjective in the case that the action is locally transitive. In this way the Mackey quantisation results above can in principle induce quantisations of T^*N .

In terms of functions on T^*N , $f \in C^\infty(N)$ corresponds to $\bar{f} = \pi^* f \in C^\infty(T^*N)$ and $v \in \mathfrak{g}$ corresponds to the function $\bar{v}(x, p) = \langle p, v(x) \rangle$. The elements of the algebra we order putting all elements of $\mathfrak{g}$ to the right. We get the relation $\bar{v} \bar{f} = \hbar v \triangleright \bar{f} + \bar{f} \bar{v}$. In terms of commutators, $[\bar{v}, \bar{f}] = \hbar v \triangleright \bar{f}$, and we would like this to be given by a Poisson bracket on T^*N . This means $\omega(d\bar{v}, d\bar{f}) = \overline{v \triangleright f}$, or in (x, p) coordinates

$$\begin{aligned} \frac{\partial \bar{f}}{\partial x^j} v(x)^j &= \omega^{(i+n)j} \frac{\partial \bar{v}}{\partial p_i} \frac{\partial \bar{f}}{\partial x^j} + \omega^{ij} \frac{\partial \bar{v}}{\partial x^i} \frac{\partial \bar{f}}{\partial x^j} \\ &= \omega^{(i+n)j} v(x)^i \frac{\partial \bar{f}}{\partial x^j} + \omega^{ij} p_k \frac{\partial v^k}{\partial x^i} \frac{\partial \bar{f}}{\partial x^j}. \end{aligned}$$

Provided the vector fields $v(x)$ for the action of $\mathfrak{g}$ span the tangent space at each point, this implies that $\omega^{(j+n)j} = 1 = -\omega^{j(j+n)}$ and all others are zero, i.e. the standard symplectic form on T^*N . Now we calculate

$$\omega(d\bar{v}, d\bar{w}) = v^j p_k \frac{\partial w^k}{\partial x^j} - w^j p_k \frac{\partial v^k}{\partial x^j} = \overline{[v, w]}.$$

These conditions are met for T^*G with Mackey quantisation $C^\infty(G) \rtimes U_\hbar(\mathfrak{g})$ where the action is by left translation. Hence in this case the formulae in Section 6.1 define an actual compatible connection ∇ on T^*G . This model will be investigated further elsewhere.

7. QUANTUM GROUPS $\mathbb{C}_q[G]$ AND RELATED EXAMPLES

Here, for completeness, we show that Drinfelds original construction of quantum groups $\mathbb{C}_q[G]$ (this means more precisely the dual of Drinfeld's construction) can also be formulated as a cochain module twist. This is not fundamentally new but a useful point of view that motivated the above. Actually, this is a general observation for any twist, in the setting of Section 2. Thus, if we are interested in A an initial undeformed Hopf algebra, let A' be a dually paired Hopf algebra and $H = A' \otimes A'^{\text{op}}$. The use of a dual here is of course avoided if we work with comodule twists rather than module ones. In our case A' acts on A from the left by $h \triangleright a = (\text{id} \otimes h)(\Delta a)$ and A'^{op} acts by $h \triangleright a = (h \otimes \text{id})(\Delta a)$. In this way A becomes H -covariant as an algebra. Now let $f \in A' \otimes A'$ be a cochain. This induces a cochain

$$F = f_{13} f_{24}^{-1} \in H \otimes H$$

where the suffices refer to the position in the four-fold tensor power of A' . The modified algebra A_F induced by this cochain has product

$$a \bullet b = \cdot(f_{24} f_{13}^{-1} \triangleright (a \otimes b)) = f(a_{(1)}, b_{(1)}) a_{(2)} b_{(2)} f^{-1}(a_{(3)}, b_{(3)})$$

which from the Drinfeld point of view is a usual (co)twist of the Hopf algebra A into a coquasiHopf algebra. Moreover, A_F will be covariant under H_F with coproduct

$$\Delta_F(h \otimes g) = f_{13} f_{24}^{-1} \Delta_{13}(h) \Delta_{24}(g) f_{24} f_{13}^{-1}$$

where the products are in the square of $A' \otimes A'^{\text{op}}$. If we denote by A'_f the usual twist of A' by f with coproduct $\Delta_f(h) = f(\Delta h) f^{-1}$. We see that $H_F = A'_f \otimes (A'_f)^{\text{op}}$, all as potentially quasi-Hopf algebras. This is the general situation and amounts to a recasting of the standard Drinfeld (co)twist of A as a module-algebra cochain twist for a suitably doubled up H .

We can of course apply this to Drinfelds example where $A = \mathbb{C}[G]$ is a suitable form of the coordinate ring of a classical simple Lie group with Lie algebra $\mathfrak{g}$, and $A' = U(\mathfrak{g})$. It is understood that we extend the above to allow formal powerseries in a parameter $\hbar$ (with $q = e^{\frac{\hbar}{2}}$). In this case Drinfeld showed the existence (as a formal power-series) of a suitable f such that $A'_f \cong U_q(\mathfrak{g})$ as a Hopf algebra. Here the coboundary ϕ_f and hence the associator Φ_f are nontrivial but ϕ_f is central in the sense that A'_f remains an ordinary Hopf algebra. This translates in the above reformulation into the statement that F and its coboundary $\phi_F \in H \otimes H \otimes H$ are nontrivial but that ϕ_F acts trivially on A , so that $A_F \cong \mathbb{C}_q[G]$ remains associative, namely the usual quantum group coordinate algebra. We have $\mathcal{L} = \mathfrak{g} \oplus \mathfrak{g}^{\text{op}}$ as the including Lie algebra from our current point of view. We can further construct $\Omega(\mathbb{C}_q[G]) = \Omega(G)_F$ and compute its associated preconnection and curvature (which is nonzero). These results are the same as those presented in [BM1] (albeit from a different supercoquasiHopf algebra point of view) so we do not repeat them here.

Note that Drinfeld's f is not known very explicitly (except at lowest order where the Poisson-bracket induced by the above is the usual Drinfeld-Sklyanin one), however its existence holds very generally and in a canonical way for quantum-group related examples. Moreover, it can happen that $H = U_q(\mathfrak{g})$ and $F = f$ may again have ϕ_F acting trivially on a particular classical algebra,

such as on a highest weight orbit. Here $\mathcal{L} = \mathfrak{g}$ is the inducing Lie algebra and we use Drinfeld's cochain without any doubling. This was the case in [DGM] where it was an associative quantum sphere was constructed in this way with $\mathcal{L} = su_2$. This is therefore an early example of the cochain-quantisation method genuinely used.

8. HIDDEN NONASSOCIATIVITY

In the above we have recovered, at least to some order, several standard associative quantum algebras of interest in physics as cochain twists (we do not just mean q-deformed or quantum group examples). Here the cochains are not required to be cocycles and this relaxation appears to be necessary. It means, however, that even though the algebra of 'functions' happens to remain associative, there is an underlying nonassociativity behind the scenes in all these quantum algebras. We now turn to this aspect.

First of all, as our algebras become quantized, their covariance Lie algebra $\mathcal{L}$ gets deformed to a quasi-quantum group $U(\mathcal{L})_F$ as explained in Section 2. These are in principle 'noncoassociative' and are looked at for our various examples in Section 8.1. Next, our quantum algebras are all equivalent in a certain monoidal categorical sense to the unquantised algebras, with the result that not only the algebras but all functorial constructions on them are similarly quantised, for example differential forms on the classical phase space and the Dirac operator deform naturally to the quantum algebras, but nonassociatively. We consider these in Sections 8.2 and 8.3 respectively.

8.1. The quasiHopf algebras $U(\mathcal{L})_F$. From Section 2, the deformed algebra A_F remains covariant, but under the quasi-Hopf algebra H_F . In our cases of interest $H = U(\mathcal{L})$ which is also the algebra of H_F . Its coproduct, however, is modified to

$$\Delta_F(X) = F(X \otimes 1 + 1 \otimes X)F^{-1} = X \otimes 1 + 1 \otimes X - \hbar[G^{(1)}, X \otimes 1 + 1 \otimes X] + O(\hbar^2)$$

for any $X \in \mathcal{L}$. The leading order data here defines a quasi-Lie bialgebra $(\mathcal{L}, \delta, \psi)$ where

$$\delta X = [X \otimes 1 + 1 \otimes X, G^{(1)}] = \text{ad}_X(G^{(1)}) \in \mathcal{L} \otimes \mathcal{L}.$$

If it happens that δ obeys the cojacobi identity

$$(\delta \otimes \text{id})\delta X + \text{cyclic} = 0$$

then we have an ordinary Lie bialgebra, which means that at lowest order at least, $U(\mathcal{L})_F$ remains an ordinary (not quasi) Hopf algebra. This is already the case for the Drinfeld twist examples in Section 7 and indeed the $O(\hbar^2)$ part ψ of ϕ is a multiple of the ad-invariant Cartan tensor $n \in \Lambda^3(\mathfrak{g})$ defined by the Killing form. In general the cojacobiator above is given by $\text{ad}_X(\psi)$ and this is the fundamental reason why the covariance algebra remains (co)associative for such examples. But let us see how the situation fares for our non-quantum group examples.

Thus, ψ is given for the $\mathcal{L} = so(1, 3)$ example in Section 4.5 and from the expression there one may readily compute that

$$\text{ad}_{X_i}(\psi) \neq 0, \quad \text{ad}_{Y_i}(\psi) = 0$$

i.e. rotationally invariant (as to be expected as the whole construction is) but not invariant under boosts. Thus our 'sphere at infinity' example in Section 4 gives us a quasi-Hopf algebra version of $\mathcal{L} = so(1, 3)$.

Next up, we CBH or $U_\hbar(g)$ example in Section 5, we have seen in Section 5.4 what the twisted coproduct Δ_F looks like to $O(\hbar^3)$ when acting on $\mathbb{C}[\mathfrak{g}] \subset H = U(\mathfrak{g} \ltimes \mathfrak{g}^*) = U(\mathfrak{g}) \bowtie \mathbb{C}[\mathfrak{g}]$. We have seen that this twists to a local form of the classical coordinate ring $\mathbb{C}[G]$. On the other hand the coproduct of $U(\mathfrak{g})$ remains unchanged after twisting to this order because all elements are g -invariant under commutator in the bigger algebra and hence

$$[\Delta v, G^{(i)}] = [v \otimes 1 + 1 \otimes v, G^{(i)}] = [v, G_1^{(i)}] \otimes G_2^{(i)} + G_1^{(i)} \otimes [v, G_2^{(i)}] = 0.$$

This is clear since the expressions involve only paired bases and dual basis, the Killing form etc. and the commutators of $v \in \mathfrak{g}$ are given by the adjoint and coadjoint actions. Such invariance would be a reasonable requirement to all orders for any universal formula for F, F^{-1} . Since $H = U(\mathfrak{g}) \bowtie \mathbb{C}[\mathfrak{g}]$ is generated by $U(\mathfrak{g}), \mathbb{C}[\mathfrak{g}]$ and has the same algebra after twisting, we conclude that

$$U(g \bowtie g^*)_F = U(g) \bowtie \mathbb{C}[g]_F$$

as an ordinary Hopf algebra. Moreover, this is locally isomorphic to $U(g) \bowtie \mathbb{C}[G] = D(U(g))$, the Drinfeld quantum double, which is an ordinary Hopf algebra. It is known that $U_{\hbar}(\mathfrak{g})$ is always covariant under $D(U(\mathfrak{g}))$ and this was explored for $U_{\hbar}(su_2)$ (the so-called universal ‘fuzzy sphere’) in [BaMa]. In this case the background covariance becomes a quantum group covariance but remains associative (there is still hidden nonassociativity, see below).

Finally, the Mackey case is a nontrivial extension of the CBH case and has a larger symmetry group. We do not make the full analysis here but suffice it to say that one reason that the CBH case works from an algebraic point of view to the fact that $U_{\hbar}(\mathfrak{g})$ is a (cocommutative) Hopf algebra and therefore has a larger quantum group covariance based on the Drinfeld double; there is no such argument for the general Mackey case but there are cases when N is itself a group and acts back on G such that the Mackey quantisation in an algebraic form becomes a bicrossproduct Hopf algebra, see [M2]. In such cases one might expect similar behaviour to the CBH case above, but not in general. The bicrossproduct case includes the deformed Poincaré group for the noncommutative spacetimes mentioned in Section 5.6, see [MR].

8.2. Quasiassociative quantum differential calculi. Next, by applying the same cochain twist to the classical exterior algebra, we obtain noncommutative differential calculi on our various quantisations. We mean differential calculus in the sense of noncommutative geometry but in a monoidal weakly associative category, and we will see that our calculi on these examples are indeed nonassociative. They do, however, have the merit of classical dimensions in each degree. Again, the quantum group case in Section 7 was already covered in an equivalent form in [BM1] with the main result that the resulting $\Omega(\mathbb{C}_q[G])$ have curvature and hence are not associative even though the quantised algebra happens to be. But let us see how our non-quantum group-related examples fare.

First, for our sphere at infinity example. The given vector fields act by Lie derivative, which commutes with the d operator. Some large calculations give the special cases:

$$\begin{aligned} x \bullet dx &= x \cdot dx + \frac{\hbar}{2z} (x^2 y \cdot dx + x(1 - x^2) \cdot dy) + \frac{\hbar^2}{8} x \cdot dx + O(\hbar^3), \\ x \bullet dy &= x \cdot dy + \frac{\hbar}{2z} (xy^2 \cdot dx + y(1 - x^2) \cdot dy) + \frac{\hbar^2}{8} x \cdot dy + O(\hbar^3), \\ y \bullet dx &= y \cdot dx - \frac{\hbar}{2z} (x(1 - y^2) \cdot dx + x^2 y \cdot dy) + \frac{\hbar^2}{8} y \cdot dx + O(\hbar^3), \\ y \bullet dy &= y \cdot dy - \frac{\hbar}{2z} (y(1 - y^2) \cdot dx + xy^2 \cdot dy) + \frac{\hbar^2}{8} y \cdot dy + O(\hbar^3). \end{aligned}$$

Moreover, since the connection ∇ arising from the noncommutativity of the calculus at lowest degree turned out to be the Levi-Civita one, and since this has (constant) curvature, we know that the exterior algebra of this quantised sphere is necessarily nonassociative.

Next, for the CBH or $U_{\hbar}(g)$ example, again we have found expressions for the curvature in terms of a double commutators. Whether or not this vanishes depends on the Lie algebra in question: for the Heisenberg Lie algebra for example, one has $R = 0$. However, for a simple Lie algebra such double commutators will not vanish and there is curvature, hence nonassociativity of $\Omega(U_{\hbar}(\mathfrak{g}))$. One can also write the deformed calculus explicitly:

$$v \bullet dw = vdw + \hbar \beta(e^i \triangleright v) de_i \triangleright w + O(\hbar^3) = vdw + \hbar \beta d[v, w] + O(\hbar^3)$$

$$dw \bullet v = (dw)v + \hbar \alpha (de_i \triangleright w) e^i \triangleright v + O(\hbar^3) = (dw)v + \hbar \alpha d[v, w] + O(\hbar^3)$$

where we use the expression for F^{-1} in Section 5.4. The second order terms fail to contribute because $e^i \triangleright v = v^i \cdot 1$ is degree zero (it acts by differentiation) which is then killed by d . Therefore the only term that could contribute in the first line, for example, is from Q_2^2 i.e. $(e^i e^j \triangleright v) de_i e_j \triangleright w$ which is zero because of the second differentiation on v . The difference between the two expressions is of course

$$v \bullet dw - dw \bullet v = \frac{\hbar}{2} d[v, w] + O(\hbar^3),$$

i.e. the Poisson-compatible preconnection as it should be. The example of ‘noncommutative spacetime’ with nonassociative differentials

$$t \bullet dx_i - dx_i \bullet t = dt \bullet x_i - x_i \bullet dt = \frac{\hbar}{2} dx_i$$

is very different from the associative (but not canonical) differential calculus usually used for this model. It represents a different approach that may overcome some of the structural problems encountered previously (such as to find the canonical Dirac operator, see below). Note that in the physical application the deformation parameter that we have denoted $\hbar$ should be denoted by another symbol and is expected to be of the order of the Planck time $\sim 10^{-44}$ s.

Finally, the Mackey quantisation case is more complicated but from the curvature computations in Section 5 we conclude again that the natural deformed calculus $\Omega(C^\infty(N) \rtimes U_\hbar(\mathfrak{g}))$ is nonassociative at least for simple Lie algebras, because the CBH part is. It is interesting to note that the curvature comes from this part alone.

8.3. Isospectral quantum Dirac operator. Here we conclude with an example to demonstrate that the categorical deformation method outlined in Section 2 is very powerful indeed and quantizes almost any natural construction. In other words, when a quantisation is expressed as a cochain module algebra twist this has great consequences.

Specifically, in another approach to noncommutative geometry it is normal to look for an analogue of the Dirac operator in the form of a ‘spectral triple’ $[C]$ obeying some axioms. These axioms are natural from an associative point of view but it is well known that important examples such as $\mathbb{C}_q[G]$ do not admit operators obeying exactly those axioms. We see by contrast that there is a natural deformation of any classical Dirac operator on the classical phase space but it will obey a variation of Connes axioms due to the hidden nonassociativity in the underlying differential calculus and elsewhere. We explain now that such an approach agrees with recent ‘isospectral deformation’ proposal for the Dirac operator on $\mathbb{C}_q[SU_2]$ in [DLSSV]. On the other hand, it is more categorical and works in principle for all quantum groups $\mathbb{C}_q[G]$, and moreover works for our more conventional quantisations such as $U_\hbar(g)$ and the Mackey quantisation to provide (in principle at least) some type of Dirac operators on them.

We consider for the sake of discussion only the case where the classical and hence quantum cotangent and spin bundles are trivial so that the spin bundle in particular has the form $V \otimes A$ where $A = C^\infty(M)$ and V is ostensibly the representation space for the spin group. We do require everything to be covariant under a background Lie algebra $\mathcal{L}$ (or Hopf algebra H) to induce the quantisation given a cochain. This is not a problem in the case $M = G$ a Lie group (a covariant Dirac operator). The short version of the quantisation is then as follows: we consider the classical Dirac operator $D : V \otimes A \rightarrow V \otimes A$ and to this we apply the functor $\mathcal{T}$ in Section 2 to obtain a map $\mathcal{T}(D) : \mathcal{T}(V \otimes A) \rightarrow \mathcal{T}(V \otimes A)$. As in Section 2 we have to allow that although $\mathcal{T}$ acts as the identity on objects and morphisms (so $\mathcal{T}(A) = A$ which becomes the deformed algebra A_F , $\mathcal{T}(V) = V$, $\mathcal{T}(D) = D$), it is nontrivial as a monoidal functor and in the sense of potentially nontrivial natural isomorphisms $\mathcal{T}(V) \otimes \mathcal{T}(A) \cong \mathcal{T}(V \otimes A)$ with certain properties in relation to $\otimes$.

We refer to [M2] for an introduction. Here these isomorphisms are given by the action of F^{-1} . Putting these facts together, we have the deformed Dirac operator:

$$D_{\bullet} : A_F \otimes V \rightarrow A_F \otimes V, \quad D_{\bullet}(a \otimes v) = F \triangleright D(F^{-1} \triangleright (a \otimes v)).$$

The main thing to note about this construction is that since it is given by conjugation by F as an operator, it does not change the spectrum in the Hilbert space setting. It should be remarked that one would still need a lot of analysis to make these remarks fully precise.

The only other subtlety is to identify A_F (which is the same vector space as A with the deformed product) explicitly as $\mathbb{C}_q[G]$ in the case $A = \mathbb{C}[G]$. This is not trivial but we note that $\mathcal{T}$ respects sums so if one has made a Peter-Weyl decomposition of A into a direct sum of matrix algebras (as is possible for $A = \mathbb{C}[G]$ for simple G) and likewise decompose $\mathbb{C}_q[G]$ in its Peter-Weyl decomposition, we can identify the summands as matrix coalgebras. This is our interpretation of the proposed Dirac operator in [DLSSV]. On the other hand, $D_{\bullet}$ lives in a nontrivial monoidal category and has properties in which the non-associativity of the category will surely play a role. It is known that the axioms in [C] are not satisfied and we would propose to replace them by ones that take this hidden nonassociativity into account.

Finally, the longer answer to the deformed Dirac operator here is to 'get inside' its construction. One can do this too in principle as we outline now. Thus, we break D into a series of morphisms all covariant under our background Hopf algebra. We also suppose for the sake of discussion that M is parallelizable (e.g. M a Lie group) so that its (covariant) differential calculus has the form $\Omega^1(M) = A \otimes \Lambda^1$ as a (trivial) bundle associated to Λ^1 . We write $da = (\partial^i a) \tau_i$ where τ_i are a basis of Λ^1 and take this as a definition of the partial derivatives. Finally, we assume ' γ -matrices' of some form $\gamma : \Lambda^1 \otimes V \rightarrow V$ so that

$$D(a \otimes v) = \partial^i(a) \otimes \gamma_i(v) = (\text{id} \otimes \gamma)(d \otimes \text{id})(a \otimes v)$$

expresses D as a composition of morphisms (here $\gamma_i = \gamma(\tau_i \otimes ())$ would be the more conventional point of view). Note that the defining relations among the γ also needs to be invariant under the background symmetry, e.g. by an invariant metric.

We now define $\gamma_{\bullet}(\tau \otimes v) = \gamma(F^{-1} \triangleright (\tau \otimes v))$ for all $\tau \in \Lambda^1$, $v \in V$ by the same reasoning as above, i.e. the functor $\mathcal{T}$. Likewise we have $d_{\bullet} = F \triangleright d$ when $d : A \rightarrow A \otimes \Lambda^1$. Note that in the above we have not deformed d and indeed this is not deformed if we consider it to Ω^1 and identify $\mathcal{T}(\Omega^1) = \Omega^1$ in the deformed theory, $d_{\bullet}$ is a slightly different object. This now expresses the above as

$$D_{\bullet} = F D F^{-1} = (\text{id} \otimes \gamma_{\bullet}) \Phi_{A, \Lambda^1, V}(d_{\bullet} \otimes \text{id})$$

in view of the diagram:

$$\begin{array}{ccccc} \mathcal{T}(A \otimes V) & \longrightarrow & \mathcal{T}(A) \otimes \mathcal{T}(V) & & \\ \downarrow d & & \downarrow d & \searrow d_{\bullet} & \\ \mathcal{T}((A \otimes \Lambda^1) \otimes V) & \longrightarrow & \mathcal{T}(A \otimes \Lambda^1) \otimes \mathcal{T}(V) & \longrightarrow & ((\mathcal{T}(A) \otimes \mathcal{T}(\Lambda^1)) \otimes \mathcal{T}(V)) \\ \parallel & & & & \downarrow \Phi_{A, \Lambda^1, V} \\ \mathcal{T}(A \otimes (\Lambda^1 \otimes V)) & \longrightarrow & \mathcal{T}(A) \otimes \mathcal{T}(\Lambda^1 \otimes V) & \longrightarrow & (\mathcal{T}(A) \otimes (\mathcal{T}(\Lambda^1) \otimes \mathcal{T}(V))) \\ \downarrow \gamma & & \downarrow \gamma & \swarrow \gamma_{\bullet} & \\ \mathcal{T}(A \otimes V) & \longrightarrow & \mathcal{T}(A) \otimes \mathcal{T}(V) & & \end{array}$$

The large middle cell here commutes by definition of a monoidal functor (the associator in the initial category of H -covariant objects is trivial). The upper and lower left cells commute because of operations on different spaces. The upper and lower right cells commute by the definitions of $d_{\bullet}$ and $\gamma_{\bullet}$ respectively. The horizontal arrows are all given by the action of F . The vertical composition on the left is $\mathcal{T}(D) = D$, while the vertical composition on the right is the definition of $D_{\bullet}$.

As to the $\gamma_\bullet$, one should write the defining relations of γ as commuting diagrams, apply the functor $\mathcal{T}$ to obtain commuting diagrams in the deformed category, and use F to interpret them as $\gamma_\bullet$ relations in a similar manner to the above. The result is, for example, a deformed set of Clifford relations involving now Φ and the deformation of the flip map induced by F (as a symmetry in the category). The actual relations would depend on the classical set up which need not be the usual classical Clifford relations if the frame group is not the usual one (e.g. one may use a Lie group to frame itself).

REFERENCES

- [AM1] H. Albuquerque & S. Majid. Quasialgebra structure of the octonions. *J. Algebra* 220:188–224, 1999.
- [AM2] E. Akrami & S. Majid. Braided cyclic cohomology and nonassociative geometry. *J. Math. Phys.* 45:3883–3911, 2005.
- [BaMa] E. Batista & S. Majid. Noncommutative geometry of angular momentum space $U(su_2)$. *J. Math. Phys.* 44: 107–137, 2003.
- [BM1] E.J. Beggs & S. Majid. Semiclassical differential structures. math.QA/0306273, to appear *Pac. J. Math.*
- [C] A. Connes. *Noncommutative Geometry*. Academic Press, 1994.
- [DLSSV] L. Dabrowski, G. Landi, A. Sitarz, W. Suijlekom & J.C. Varilly The Dirac operator on $SU_q(2)$. Preprint, math.QA/0411609.
- [DGM] J. Donin, D. Gurevich & S. Majid. R-matrix Brackets and their Quantization. *Ann. Inst. H. Poincaré*, 58: 235–246, 1993.
- [D1] V.G. Drinfeld. Quantum groups. In A. Gleason, editor, *Proceedings of the ICM*. Pages 798–820, AMS, Rhode Island, 1987.
- [D2] V.G. Drinfeld. Quasi-Hopf algebras. *Leningrad Math. J.*, 1:1419–1457, 1990.
- [Du] M. Duflo Caracteres des algèbres de Lie résolubles. *C. R. Acad. Sci.* 269:437438, 1969.
- [F] B. Fedosov. *Deformation quantization and index theory*. Akademie Verlag, Berlin, 1996.
- [GRS] I. Gelfand, V. Retakh & M. Shubin. Fedosov manifolds. *Adv. Math.* 136:104–140, 1998.
- [G] S. Gutt. An explicit $*$ product on the cotangent bundle of a Lie group. *Lett. in Math. Phys.* 7:249–258, 1983.
- [H] E. Hawkins. Noncommutative Rigidity. *Commun. Math. Phys.* 246: 211–235, 2004.
- [M1] S. Majid. Tannaka-Krein Theorem For QuasiHopf Algebras and Other Results. *Contemp. Math.* 134:219–232, 1992.
- [M2] S. Majid. *Foundations of Quantum Group Theory*. Cambridge Univ. Press, 1995.
- [M3] S. Majid. Gauge theory on nonassociative spaces. *J. Math. Phys.* 46:103519, 23pp., 2005.
- [M4] S. Majid. Noncommutative model with spontaneous time generation and Planckian bound. *J. Math. Phys.* 46:103520, 18pp., 2005.
- [MO] S. Majid and R. Oeckl. Twisting of quantum differentials and the Planck scale Hopf algebra. *Commun. Math. Phys.*, 205:617–655, 1999.
- [MR] S. Majid and H. Ruegg. Bicrossproduct structure of the k -Poincaré group and non-commutative geometry. *Phys. Lett. B*, 334:348–354, 1994.

EJB: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WALES, SWANSEA SA2 8PP, UK, SM: SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY, UNIVERSITY OF LONDON, 327 MILE END RD, LONDON E1 4NS, UK